arXiv:1209.0881v1 [math-ph] 5 Sep 2012

The Physics of Events: A Potential Foundation for Emergent Space-Time

Kevin H. Knuth
Departments of Physics and Informatics
University at Albany (SUNY)
Albany NY 12222, USA

Newshaw Bahreyni
Department of Physics
University at Albany (SUNY)
Albany NY 12222, USA

September 6, 2012

Abstract

Everything that is detected or measured is the direct result of something influencing something else. This is the essence of the concept of force, which has become central to physics. By considering both the act of influencing and the response to such influence as a pair of events, we can describe a universe of interactions as a partially-ordered set of events. In this paper, we take the partially-ordered set of events as a fundamental picture of influence and aim to determine what interesting physics can be recovered. This is accomplished by identifying a means by which events in a partially-ordered set can be aptly and consistently *quantified*. Since, in general, a partially-ordered set lacks symmetries to constraint any quantification, we propose to distinguish a chain of events, which represents an observer, and quantify some subset of events with respect to the observer chain. We demonstrate that consistent quantification with respect to pairs of observer chains exhibiting a constant relationship with one another results in a metric analogous to the Minkowski metric and that transformation of the quantification with respect to one pair of chains to quantification with respect to another pair of chains results in the Bondi k-calculus, which represents a Lorentz transformation under a simple change of variables. We further demonstrate that chain projection induces geometric structure in the partially-ordered set, which itself is inherently both non-geometric and non-dimensional. Collectively, these results suggest that the concept of space-time geometry may emerge as a unique way for an embedded observer to aptly and consistently quantify a partially-ordered set of events. In addition to having potential implications for space-time physics, this also may serve as a foundation for understanding analogous space-time in condensed matter systems.

1 Introduction

Everything that is detected or measured is the direct result of something influencing something else. This is the essence of the concept of force, which has become central to physics. When it is possible to clearly define an entity, such as a stable particle, one can talk of particles influencing one another as in the most common conception of force. One can also conceive of a particle influencing itself, which can be thought of in terms of a force mediating some kind of self-reaction resulting in dynamic evolution. Influence is central also in cases where entities cannot be well-delineated, such as in particle decays. The common feature of each of these situations is that something influences and something is influenced.

Here we focus on an admittedly simplistic, but fundamental, picture of influence where we consider the process of influence to connect and order the act of influencing and the act of being influenced. We refer to each of these two acts with the generic term ***event***, so that the event associated with the act of influencing influences the event associated with the act of being influenced. This generic process of influence, along with the notion of transitivity of such influence, allows events to be partially-ordered. This results in a structure referred to mathematically as a ***partially-ordered set***, or a ***poset*** for short. In this exploration, we will not be concerned with differentiating between distinct types of influence and their corresponding distinct types of events. Instead, with the aim toward illuminating the foundations, the goal here is to determine what interesting physics can be recovered from this minimalist picture.

Partially-ordered sets of events ordered by causal influence were introduced by Bombelli et al. [3] and called ***causal sets*** or ***causets***. Over the last twenty years, causal sets have been championed by Rafael Sorkin [27, 28] and employed in approaches to quantum gravity. As such they are typically endowed with, or embedded within, a Minkowski geometry exhibiting Lorentz invariance [4]. We approach the problem from another direction entirely. We take the partially-ordered set of events as a fundamental picture of influence. Rather than endowing it with additional properties, our goal is simply to identify a consistent means by which events in the poset can be aptly ***quantified***. Where assumptions or hypotheses regarding the properties of events or relationships between events have the potential to be right or wrong, consistent apt quantification can only be useful or not useful.

From our previous studies in quantifying lattices, which are special cases of partially-ordered sets[1] [16, 17, 18, 19, 20, 23], we have found that the underlying order constrains quantification. Most recently, we have demonstrated how the algebraic (equivalently order-theoretic) relations between experimental setups constrain apt quantification of measurement sequences resulting in the complex sum and product rules [12, 11] of the Feynman path integral formulation of quantum mechanics [9]. In contrast to the topics of previous study, a partially-ordered set of events has no symmetry in general that can constrain a quantification scheme. In this paper, we aim to determine what minimal structure is necessary to obtain useful constraints on the quantification of partially-ordered sets of events, and to determine precisely what those constraints are.

---

[1] Lattices are partially-ordered sets where every pair of elements has a unique least upper bound called a join and a greatest lower bound called a meet. This enables one to consider the join and meet as algebraic operators so that all lattices are algebras.

In the next section we introduce the concept of a distinguished chain, which can be used to represent an observer. Quantification with respect to distinguished chains is explored in Section 3 and extended to the concept of intervals between events in Section 4. There we demonstrate that consistent quantification with respect to pairs of chains exhibiting a constant relationship with one another results in a metric analogous to the Minkowski metric and that transformation of the quantification with respect to one pair of chains to quantification with respect to another pair of chains results in the Bondi k-calculus [5], which represents a Lorentz transformation under a simple change of variables. It is important to keep in mind that this is obtained within the framework of a partially-ordered set where there is no fundamental concept of motion, and thus no opportunity to assume that there exists a constant and maximal speed of light. Instead, these relations arise to enforce the partial-order imposed by the concept of influence. Section 5 explores the structure introduced by describing the poset with respect to an embedded observer chain. Such structure includes the concept of induced subspaces, which leads to a natural concept of dimension along with many other "geometric" concepts including a novel derivation of the Pythagorean theorem. Collectively, these results suggest that the concept of space-time geometry emerges as the unique way for an embedded observer to aptly and consistently quantify a partially-ordered set of events.

2 Events, Chains and Observers

Influence is bounded by two events: the action of influencing and the reaction of being influenced. As such, influence can be viewed as a binary ordering relation, which relates pairs of events. That is, if event x influences event y, we write $x \leq y$, and generically read 'y ***includes*** x'. We take as a postulate that influence is transitive. This ensures that this binary ordering relation imposes a partial order on the set of events so that if $x \leq y$ and $y \leq z$, then it is also true that $x \leq z$. Given any pair of events, it is not always true that one event is influenced by the other. In this case, we say that the events are ***incomparable*** and write $x \,||\, z$. We further define the binary ordering relation so that $x \leq y$ and $y \leq x$ can hold simultaneously only if $x = y$. Taken together, these postulates enable one to describe events in terms of a ***partially-ordered set***, or ***poset*** for short.

Some posets, such as lattices, possess symmetries that give rise to algebraic structures that can be used to guide consistent quantification [16, 19, 23, 12, 11]. However, this is not the case for posets in general where often the only structure present is the partial ordering itself. Given that our present goal is to discover what minimal structure is necessary to obtain useful constraints on the consistent apt quantification of events, we propose to introduce additional structure simply by ***distinguishing*** a set of events, such as a finite chain, and quantifying a subset of the poset with respect to the distinguished set.

A chain is a subset of poset elements where for every element x and y in the chain we have that either $x \leq y$ or $y \leq x$ so that the elements comprising the chain are totally ordered. In other words, a chain consists of a set of events which occur in succession. Physically, such a chain of events may represent the ticks of a clock or more fundamentally a particle evolving in time by experiencing a sequence of detectable changes in its internal state. It is the mental image of an observer with a clock taking note of the time at which events are observed that leads us to refer to the distinguished chain as an ***observer chain***.

In the next section, we focus on mathematics and introduce a consistent means by which one can quantify elements on a chain as well as the intervals between elements on a chain. By extending such quantification to a subset of the poset of events, we obtain a set of mathematical relations that enable a quantitative description of events in general.

3 Quantification of a Poset by Chain Projection

In this section, we focus on mathematics and examine the quantification of a partially-ordered set by introducing a method which we call ***chain projection***. We limit ourselves to partially-ordered sets Π in which a finite chain $\mathbf{P}$ comprised of N elements $p_1 < p_2 < \ldots p_N \in \Pi$ can be identified, and introduce a consistent method by which the chain can be quantified. By introducing the notion of a projection onto a chain, we extend the quantification of a chain to a subset of the poset. Note that we will find that it is not guaranteed that the entire poset can be quantified in such a manner. This fact has important consequences for the resulting physics of events.

3.1 Quantification of a Chain

Given a poset Π, we begin by distinguishing a chain $\mathbf{P}$ comprised of N elements $p_1 < p_2 < \ldots p_N \in \Pi$. Quantification is introduced by means of an ***isotonic valuation***, which is a functional that assigns a real number to each element in the chain $\mathbf{P}$, $v_{\mathbf{P}} : p_x \in \mathbf{P} \to I\!\!R$, such that for any two elements $p_x, p_y \in \mathbf{P}$ related by $p_x < p_y$ we have that $v_{\mathbf{P}}(p_x) \leq v_{\mathbf{P}}(p_y)$. Note that the case where $p_x < p_y$ and $v_{\mathbf{P}}(p_x) = v_{\mathbf{P}}(p_y)$ is intentional, and amounts to a form of coarse-graining where the elements p_x and p_y are not distinguished by the selected quantification scheme. The result is that the chain $\mathbf{P} = \{p_1 < p_2 < \ldots p_N\} \in \Pi$, is quantified by a non-decreasing sequence of real numbers $\{v_{\mathbf{P}}(p_1) \leq v_{\mathbf{P}}(p_2) \leq \ldots v_{\mathbf{P}}(p_N)\}$.

3.2 Chain Projection Mapping

We consider the possible relationships between the chain $\mathbf{P}$ and elements of the poset Π. Given an element $x \in \Pi$ and $x \notin \mathbf{P}$, one of the four following situations (illustrated in Figure 1) holds:

$$A. \quad \left\{ \begin{array}{lll} p_i \,||\, x & \forall & 1 \leq i \leq N \end{array} \right. \qquad (\mathbf{P} \text{ and } x \text{ are incomparable})$$

$$B. \quad \left\{ \begin{array}{lll} p_i \leq x & \forall & 1 \leq i \leq L \\ p_i \,||\, x & \forall & L < i \leq N \end{array} \right. \qquad (x \text{ includes elements of } \mathbf{P})$$

$$C. \quad \left\{ \begin{array}{lll} p_i \,||\, x & \forall & 1 \leq i < U \\ p_i \geq x & \forall & U \leq i \leq N \end{array} \right. \qquad (\text{elements of } \mathbf{P} \text{ include } x)$$

$$D. \quad \left\{ \begin{array}{lll} p_i \leq x & \forall & 1 \leq i \leq L \\ p_i \,||\, x & \forall & L < i < U \\ p_i \geq x & \forall & U \leq i \leq N \end{array} \right. \qquad (\mathbf{P} \text{ and } x \text{ include one another})$$



Figure 1: Illustration of the four possible relationships between a chain $\mathbf{P}$ and an element $x \notin \mathbf{P}$. Note that in (D) the element x can be mapped to two elements on $\mathbf{P}$, $\overline{P}_x = p_L$ and $P_x = p_U$. Poset elements exhibiting this relationship to the chain can be quantified in two ways by inheriting the quantification $v_{\mathbf{P}}(\overline{p}_x)$ and $v_{\mathbf{P}}(p_x)$ of the elements $\overline{p}_x$ and p_x.

We say that the poset element x can be ***projected*** onto a chain $\mathbf{P}$ if there exists an element $p \in \mathbf{P}$ such that $x \leq p$ (Cases C and D). If this is the case, then the ***forward projection***, or simply ***projection***, of x onto the chain $\mathbf{P}$ is given by the least event p_x on the chain $\mathbf{P}$ such that $x \leq p_x$ where $p_x := \inf\{p \in \mathbf{P} | x \leq p\}$, which is indicated by the index U above. Since the projection of an element onto a chain, if it exists, is unique, we can consider the projection to be a functional $P : x \in \Pi \rightarrow p_x \in \mathbf{P}$. By applying this functional to the element x, we have $p_x = Px$.

One can also consider the ***dual projection***[2] or ***backward projection*** $\overline{P}$ where one identifies the greatest element $\overline{p}_x$ on the chain that is included by the poset element x, so that $\overline{p}_x = \overline{P}_x := \sup\{p \in \mathbf{P} | p \leq x\}$ corresponding to the element indexed by L above (Cases B and D).

The forward and backward projections together result in a surjective ***chain projection map***, which provides information about the connectivity of a poset

---

[2] Note that the dual projection $\overline{P}_x$ of x onto the chain $\mathbf{P}$ is equal to the projection of x onto the dual chain $\mathbf{P}^\partial$ where the order is reversed.



Figure 2: This figure illustrates the quantification of a poset with respect to the chain **P**, differentiated by the bold connections. The chain **P** is quantified by an isotonic valuation consisting of integers. Note that the repeated assignment of 4 is intentional. Poset elements, such as x, are then quantified by pairs of the form $(p_x, \overline{p}_x)$ representing the projection and the dual projection onto the chain **P**. Note that not all elements can be quantified by two numbers. Some elements, such as those quantified by $(., 3)$ do not project to the chain **P**. Others, such as $(2, .)$ do not possess a dual projection. Last, some elements, such as the one quantified by $(., .)$ are incomparable to all elements of the chain and cannot be quantified this way.

from the perspective of the observer chain. Any ***range*** of the chain projection map, which is defined by two intervals $[\overline{p}_{min}, \overline{p}_{max}]$ and $[p_{min}, p_{max}]$ where $\overline{p}_{min} < p_{max}$, implicitly defines a domain of poset elements.

By composing the chain projection map with the valuation map from chain elements to real numbers, the domain of the valuation can be extended from the chain itself to the domain corresponding to any quantified range along the chain so that for a chain **P** we have $v_{\mathbf{P}}(x) := v_{\mathbf{P}}(Px)$ and $\overline{v}_{\mathbf{P}}(x) := v_{\mathbf{P}}(\overline{P}x)$.

Given the projection and the dual projection, we can quantify elements that both include and are included by elements of the chain (Case D) with a pair of numbers. That is, given a chain **P** and element x where both Px and $\overline{P}x$ exist, we can quantify x using a pair of numbers $(v_{\mathbf{P}}(Px), v_{\mathbf{P}}(\overline{P}x))$. Figure 2 illustrates quantification of a subset of poset events by chain projection. Such quantification results in a set of ***chain-based coordinates***. Note that an element x on the chain **P** quantified by the scalar $v_{\mathbf{P}}(x)$ is also quantified by the pair $(v_{\mathbf{P}}(x), v_{\mathbf{P}}(x))$ since $x = Px = \overline{P}x$ for every $x \in \mathbf{P}$.

While this quantification scheme does not ensure that all elements in the



Figure 3: This figure illustrates the lattice of partitions of closed intervals defined on a chain. The join operation on this lattice is represented by the operator $\cup$ that joins pairs of closed intervals that have elements in common, such as $[1,3] \cup [2,4] = [1,4]$. The meet operation is given by set intersection $\cap$.

partially-ordered set will be quantified, we will find it to be extremely useful.

4 Intervals

Often we are interested in the relationship between two elements. In this section we define the concept of closed intervals on a chain, which will later be extended to the concept of a generalized interval that may be defined by considering incomparable elements.

4.1 Closed Intervals

Given two elements $p_a \leq p_b \in \mathbf{P}$ one can define a ***closed interval*** $[p_a, p_b]_\mathbf{P}$ as the set of elements $x \in \mathbf{P}$ such that $p_a \leq x \leq p_b$ [2, p.7].[3] The idea is that the closed interval $[p_a, p_b]_\mathbf{P}$ indicates a relationship between the two elements on the chain given by the boundary of the interval $\{p_a, p_b\} = \partial[p_a, p_b]_\mathbf{P}$.

Two closed intervals on a chain $\mathbf{P}$ that have elements in common can be combined to form a single closed interval. For example, given the intervals $[a, c] = \{a, b, c\}$ and $[b, d] = \{b, c, d\}$ we have that

$$[a, c] \cup [b, d] = [a, d], \tag{1}$$

where $\cup$ is the usual set union operation so that $[a, d] = \{a, b, c, d\}$. Note that we have dropped the subscript indicating that the elements comprising this interval belong to the chain $\mathbf{P}$. Closed intervals themselves can be ranked based on whether one closed interval is a subset of another resulting in a distributive lattice. An example of such a lattice is shown in Figure 3 for the case of integral partitions of the interval $[1, 5]$. These structures are closely related to Young diagrams or Young tableaux, which are another way of representing partitions of an integer.

Ranking can be generalized to quantification by defining a valuation ϕ that takes a closed interval $I_\mathbf{P}$ to a real number, $\phi : I_\mathbf{P} \rightarrow I\!R$, such that when the closed interval $I_{\mathbf{P}}$ is a subset of the closed interval $J_{\mathbf{P}}$, $I_{\mathbf{P}} \subseteq J_{\mathbf{P}}$, we have that $\phi(I_{\mathbf{P}}) \leq \phi(J_{\mathbf{P}})$. Such a quantification scheme results in a calculus of closed intervals, which is constructed to numerically encode their algebraic properties (see for example [16, 19, 12]).

---

[3] The subscript $\mathbf{P}$ in $[p_a, p_b]_\mathbf{P}$ indicates that $p_a \leq x \leq p_b \in \mathbf{P}$.

General results must apply to specific cases, so we begin by considering the special case involving the joining of pairs of closed intervals that share only a common endpoint

$$[a, b] \cup [b, c] = [a, c]. \tag{2}$$

If the valuation assigned to the intervals is to encode their relationships to one another, then it must be that $\phi([a, b] \cup [b, c])$ is a function of $\phi([a, b])$ and $\phi([b, c])$ so that

$$\phi([a, b] \cup [b, c]) = \phi([a, b]) \oplus \phi([b, c]), \tag{3}$$

where $\oplus$ is an operator (to be determined) that acts on the valuations.

We note also that join operation $\cup$ is associative

$$[a, b] \cup ([b, c] \cup [c, d]) = ([a, b] \cup [b, c]) \cup [c, d], \tag{4}$$

so that the order in which closed intervals are joined does not affect the final result. We require that the valuation ϕ encodes this property of associativity. If it did not, then one could not consistently assign a unique scalar valuation to a composite closed interval such as $[a, d]$ in the equation above since the order in which the closed intervals were combined to form the composite would affect the final valuation. For the valuation ϕ to obey associativity, we require that

$$\phi([a, c]) \oplus (\phi([b, c]) \oplus \phi([c, d])) = (\phi([a, b]) \oplus \phi([b, c])) \oplus \phi([c, d]). \tag{5}$$

If we write $\alpha = \phi([a, b])$, $\beta = \phi([b, c])$, $\gamma = \phi([c, d])$ then the equation above becomes

$$\alpha \oplus (\beta \oplus \gamma) = (\alpha \oplus \beta) \oplus \gamma, \tag{6}$$

which is a functional equation for $\oplus$ called the *associativity equation* [1, 23]. Its general solution is

$$\alpha \oplus \beta = f^{-1}(f(\alpha) + f(\beta)), \tag{7}$$

where f is an arbitrary invertible function [1, 23]. This means that without loss of generality, we can always perform a regraduation to a valuation $d = f \circ \phi$ so that $d([a, b]) = f(\phi([a, b]))$ and the valuations assigned to the appropriate closed intervals sum when they are combined as in (2)

$$d([a, c]) = d([a, b] \cup [b, c]) = d([a, b]) + d([b, c]). \tag{8}$$

It is straightforward to show that in general for any pair of closed intervals $I_{\mathbf{P}}$ and $J_{\mathbf{P}}$ sharing at least one element so that $I_{\mathbf{P}} \cap J_{\mathbf{P}} \neq \emptyset$, the valuation obeys the ***sum rule*** [16, 17, 19, 23], which is also known as the ***inclusion-exclusion relation*** [15]

$$d(I_{\mathbf{P}} \cup J_{\mathbf{P}}) = d(I_{\mathbf{P}}) + d(J_{\mathbf{P}}) - d(I_{\mathbf{P}} \cap J_{\mathbf{P}}). \tag{9}$$

This can be viewed as a constraint equation that respects both the ordering of the intervals and associativity of their join, and can be directly compared to the familiar sum rule in probability theory [16, 17, 19, 23].



Figure 4: Since the join operation that combines closed intervals is associative, $[a, b] \cup ([b, c] \cup [c, d]) = ([a, b] \cup [b, c]) \cup [c, d]$, the operation $\oplus$ representing the combination of valuations (lengths) quantifying the intervals is also associative so that $\alpha \oplus (\beta \oplus \gamma) = (\alpha \oplus \beta) \oplus \gamma$.

Since a closed interval $[a, b]$ is defined by its extreme elements a and b, we look for a relationship between the valuations assigned to these two elements and the valuation assigned to the closed interval

$$d([a, b]) = u(v(a), v(b)), \tag{10}$$

where u is a function to be determined. We can now rewrite $d([a, c]) = d([a, b]) + d([b, c])$ as

$$u(v(a), v(c)) = u(v(a), v(b)) + u(v(b), v(c)), \tag{11}$$

which has as its unique solution

$$u(v(a), v(c)) = v(c) - v(a), \tag{12}$$

so that, without loss of generality,

$$d([a, c]) = v(c) - v(a) \tag{13}$$

provides a scalar quantification of the closed interval given the selected grade and the valuations assigned to the elements of the chain. We will refer to this scalar quantification of a closed interval on a chain as the ***length*** of the interval.

In summary, a closed interval $[p_a, p_b]_{\mathbf{P}}$ defined on a chain $\mathbf{P}$ can be quantified by a pair of numbers $\big(v_{\mathbf{P}}(p_a), v_{\mathbf{P}}(p_b)\big)$ as well as a scalar $v_{\mathbf{P}}(p_b) - v_{\mathbf{P}}(p_a)$. In the next section, we generalize the notion of an interval to include incomparable elements.

4.2 Generalized Intervals

We extend the concept of a closed interval on an ordered chain to that of a ***generalized interval***. A generalized interval, denoted $[x, y]$, is identified by an ordered pair of elements $x, y \in \Pi$, each of which is called an ***endpoint***. Note that x and y are either comparable ($x \leq y$ or $y \leq x$) as in the case of a closed interval or incomparable ($x \,||\, y$) where the endpoints are elements of different chains. For this reason, the subscript referring to a chain has been dropped from the notation. Henceforth we shall refer to a generalized interval simply as an ***interval***.

The remainder of the paper will focus on deriving a quantification of intervals via chain projection.

5 Chain-Induced Structure and the Quantification of Intervals

Together the act of distinguishing an observer chain and the technique of chain projection effectively maps a subset of poset elements, and hence intervals, onto chain elements and closed intervals, respectively. This reveals information about the connectivity of the poset of events from the perspective of the distinguished chain by specifically indicating the relationship between the chain and the poset elements in the quantified subset. As such, the chain projection mapping has the effect of inducing structure in a poset that may lack any inherent characteristic structure or symmetry.

In this section, we explore the structure induced by such mappings in the case of multiple quantifying chains where closed intervals along one quantifying chain project to closed intervals along another quantifying chain. For arbitrary chains in a general poset, one does not expect there to be a relationship between such projections. However, in the special case where a set of multiple quantifying chains mutually agree in their quantification of one another we show that there exists a unique consistent pair-wise and scalar quantification of intervals. This sets the stage for an observer-based geometry.

5.1 Induced Subspaces

In this section we demonstrate how multiple chains can induce a subspace within the poset. We begin by introducing the concept of collinearity.

> ***Collinearity***: An element x is said to be ***collinear*** with finite chains $\mathbf{P}$ and $\mathbf{Q}$, denoted $x \in \overline{\mathbf{PQ}}$, iff the projections of x onto $\mathbf{P}$, Px and $\overline{P}x$, can be found by first projecting x onto $\mathbf{Q}$ and then onto $\mathbf{P}$, and vice versa by interchanging the roles of $\mathbf{P}$ and $\mathbf{Q}$.

There are three ways that this can be accomplished without violating the partial order:

$$\begin{array}{lll} Px = \overline{P}Qx & Qx = QPx & \\ \overline{P}x = P\overline{Q}x & \overline{Q}x = \overline{QP}x & \text{(Case I)} \\ & & \\ Px = PQx & Qx = \overline{Q}Px & \\ \overline{P}x = \overline{PQ}x & \overline{Q}x = Q\overline{P}x & \text{(Case II)} \\ & & \\ Px = P\overline{Q}x & Qx = Q\overline{P}x & \\ \overline{P}x = \overline{P}Qx & \overline{Q}x = \overline{Q}Px & \text{(Case III)} \end{array} \tag{14}$$

The first two cases map to one another under interchange of $\mathbf{P}$ and $\mathbf{Q}$, whereas the third case maps to itself. This result leads to a geometric interpretation of


Case I
$Qx = QPx$
$Px = \overline{P}Qx$
x
$\overline{P}x = P\overline{Q}x$
$\overline{Q}x = \overline{Q}\overline{P}x$
P
Q
Case II
$Px = PQx$
$Qx = \overline{Q}Px$
x
$\overline{Q}x = Q\overline{P}x$
$\overline{P}x = \overline{P}\overline{Q}x$
P
Q
Case III
$Px = P\overline{Q}x$
$Qx = Q\overline{P}x$
x
$\overline{P}x = \overline{P}Qx$
$\overline{Q}x = \overline{Q}Px$
P
Q

Figure 5: An illustration of the three ways in which an element x can be collinear with two finite chains. In this way, an element can be said to be either on one side or the other of, or between, a pair of chains. This introduces the concept of directionality to the subspace generated by the coordinated chains $\mathbf{P}$ and $\mathbf{Q}$.

these three relationships. To begin to understand this, the concept of collinearity enables one to divide the poset into two equivalence classes based on whether or not an element is collinear with the chains $\mathbf{P}$ and $\mathbf{Q}$. Elements that are collinear with $\mathbf{P}$ and $\mathbf{Q}$ are said to reside in the subspace defined by $\mathbf{P}$ and $\mathbf{Q}$. This subspace can be further divided into three equivalence classes according to the three cases above as illustrated in Figure 5. An element x exhibiting the relationship in Case I is said to be situated on the $\mathbf{P}$-side of the pair of chains $\mathbf{P}$ and $\mathbf{Q}$, which is denoted by $x|\mathbf{P}|\mathbf{Q}$; whereas an element x exhibiting the relationship in Case II is said to be situated on the $\mathbf{Q}$-side of the pair of chains $\mathbf{P}$ and $\mathbf{Q}$, which is denoted by $\mathbf{P}|\mathbf{Q}|x$. Last, an element x exhibiting the relationship in Case III is said to be situated *between* $\mathbf{P}$ and $\mathbf{Q}$, which is denoted by $\mathbf{P}|x|\mathbf{Q}$.



Figure 6: (Left) The element x is not only collinear with, but also between, chains $\mathbf{P}$ and $\mathbf{Q}$ since $Px = P\overline{Q}x$ and $Qx = Q\overline{P}x$. In this situation, quantification can be performed by considering only forward projections, such as Px and Qx. (Right) An example of an element x which is not collinear with the chains $\mathbf{P}$ and $\mathbf{Q}$. In this sense, the chains $\mathbf{P}$ and $\mathbf{Q}$ define a subspace, which can include or exclude elements. Note that the definition of the projection imposes the constraint that $P\overline{Q}x \leq Px$, and similarly for the other pairs of projections.

These relationships can be extended to any generalized interval $[a, b]$ resulting in nine cases, three cases where the two elements are situated similarly

$$\begin{array}{ll} [a,b]|\mathbf{P}|\mathbf{Q} & [a,b] \text{ is on the } \mathbf{P}\text{-side of } \mathbf{P} \text{ and } \mathbf{Q} \\ \mathbf{P}|\mathbf{Q}|[a,b] & [a,b] \text{ is on the } \mathbf{Q}\text{-side of } \mathbf{P} \text{ and } \mathbf{Q} \\ \mathbf{P}|[a,b]|\mathbf{Q} & [a,b] \text{ is between } \mathbf{P} \text{ and } \mathbf{Q} \end{array} \tag{15}$$

and six cases where the interval $[a, b]$ straddles one or more chains

$$\begin{array}{ll} a|\mathbf{P}|b|\mathbf{Q} & b|\mathbf{P}|a|\mathbf{Q} \\ a|\mathbf{P}|\mathbf{Q}|b & b|\mathbf{P}|\mathbf{Q}|a \\ \mathbf{P}|a|\mathbf{Q}|b & \mathbf{P}|b|\mathbf{Q}|a. \end{array} \tag{16}$$

We can extend these relationships to include finite chains. In the case where each element of a finite chain $\mathbf{X}$ exhibits the same relationship to $\mathbf{P}$ and $\mathbf{Q}$, $\mathbf{X}$ can be said to be either on the $\mathbf{P}$-side of $\mathbf{P}$ and $\mathbf{Q}$ as in $\mathbf{X}|\mathbf{P}|\mathbf{Q}$, or the $\mathbf{Q}$-side of $\mathbf{P}$ and $\mathbf{Q}$ as in $\mathbf{P}|\mathbf{Q}|\mathbf{X}$, or between $\mathbf{P}$ and $\mathbf{Q}$ as in $\mathbf{P}|\mathbf{X}|\mathbf{Q}$. This introduces a way in which some chains belonging to this subspace can be ordered, which in turn induces bi-directionality. This gives rise to an induced ordering relation where we can write

$$\mathbf{X}|\mathbf{P}|\mathbf{Q} \quad \Rightarrow \quad \begin{array}{c} \mathbf{X} < \mathbf{P} < \mathbf{Q} \\ \text{or} \\ \mathbf{Q} < \mathbf{P} < \mathbf{X} \end{array} \tag{17}$$

where the directionality of the induced ordering relation is arbitrary.

Since in this situation, we have two ordering relations: one along chains, and one among collinear chains; we say that the induced subspace is 1+1 dimensional.

It is important to note that not all poset elements are members of the subspace defined by any pair of chains as is illustrated in Figure 6(right) where the element x is said to be situated outside of the $1 + 1$ dimensional subspace induced by the coordinated chains $\mathbf{P}$ and $\mathbf{Q}$.

5.2 Coordinated Chains

If projection onto a single observer chain provides information about a poset, then one could imagine that two or more observer chains could be used to capture even more information about the poset. However, unless the relationships among the set of observer chains are known, additional information provided by the additional chains cannot be combined to provide a more detailed picture of the poset.

In this section, we consider posets that support multiple observer chains, which can be used to quantify the poset via chain projection. Furthermore, we assume that these chains are ***compatible*** and can be ***coordinated*** so that they project onto one another in a well-defined manner thus enabling us to extend the chain-based coordinate system induced by each chain to a coordinate system that is potentially more global.

> ***Compatibility***: Two chains $\mathbf{P}$ and $\mathbf{Q}$ are said to be ***compatible*** over the intervals given by $[\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$, $[p_{min}, p_{max}]_{\mathbf{P}}$ and $[\overline{q}_{min}, \overline{q}_{max}]_{\mathbf{Q}}$, $[q_{min}, q_{max}]_{\mathbf{Q}}$ iff every element $\overline{p} \in [\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$ projects to a unique element $q \in [q_{min}, q_{max}]_{\mathbf{Q}}$, and every element $\overline{q} \in [\overline{q}_{min}, \overline{q}_{max}]_{\mathbf{Q}}$ projects to a unique element $p \in [p_{min}, p_{max}]_{\mathbf{P}}$.

The concept of compatibility ensures that every sub-interval in $[\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$ on the chain $\mathbf{P}$ maps isomorphically via projection to a unique sub-interval in $[q_{min}, q_{max}]_{\mathbf{Q}}$ on the chain $\mathbf{Q}$ and vice versa. Compatibility can be extended to valuations so that the two chains agree on the quantification of the two sets of closed intervals on each chain. We refer to such a pair of chains as ***coordinated chains***.

> ***Coordinated Chains***: Two chains $\mathbf{P}$ and $\mathbf{Q}$ are said to be ***coordinated*** over the intervals given by $[\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$, $[p_{min}, p_{max}]_{\mathbf{P}}$ and $[\overline{q}_{min}, \overline{q}_{max}]_{\mathbf{Q}}$, $[q_{min}, q_{max}]_{\mathbf{Q}}$ iff they are compatible over the specified ranges and every sub-interval in $[\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$ and its projection onto a sub-interval in $[q_{min}, q_{max}]_{\mathbf{Q}}$ are quantified by the same valuation, and likewise for each sub-interval in $[\overline{q}_{min}, \overline{q}_{max}]_{\mathbf{Q}}$ and its projection into $[p_{min}, p_{max}]_{\mathbf{P}}$. That is, for $\overline{p}_x, \overline{p}_y \in [\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$, we have that $v_{\mathbf{P}}(\overline{p}_y) - v_{\mathbf{P}}(\overline{p}_x) = v_{\mathbf{Q}}(Q\overline{p}_y) - v_{\mathbf{Q}}(Q\overline{p}_x)$ for $Q\overline{p}_y, Q\overline{p}_x \in [q_{min}, q_{max}]_{\mathbf{Q}}$ and similarly for $\overline{q}_x, \overline{q}_y \in [\overline{q}_{min}, \overline{q}_{max}]_{\mathbf{Q}}$.

The concept of coordinated chains is illustrated in Fig. 7. A pair of coordinated chains $\mathbf{P}$ and $\mathbf{Q}$ generates a 1+1 dimensional subspace, denoted by $\overline{\mathbf{PQ}}$, which includes $\mathbf{P}$, $\mathbf{Q}$, and all elements collinear with $\mathbf{P}$ and $\mathbf{Q}$. In this situation, the pair of chains can be used to quantify elements between them using only forward projections, which is an advantage since in practical situations it represents information that can be obtained by the chains themselves. This is because Case III of the collinearity condition requires that $Px = P\overline{Q}x$ and $Qx = Q\overline{P}x$ so that $v_{\mathbf{P}}(\overline{P}_y) - v_{\mathbf{P}}(\overline{P}_x) = v_{\mathbf{Q}}(Q_y) - v_{\mathbf{Q}}(Q_x)$ as given by the definition of coordinated chains.



Figure 7: An illustration of two chains coordinated over the range given by $[\overline{p}_{min}, \overline{p}_{max}]_{\mathbf{P}}$, $[p_{min}, p_{max}]_{\mathbf{P}}$ and $[\overline{q}_{min}, \overline{q}_{max}]_{\mathbf{Q}}$, $[q_{min}, q_{max}]_{\mathbf{Q}}$. Any interval formed from the indicated events in between the two chains can be quantified by the same pair whether it is obtained by projection and dual projection onto **P** or **Q**, or projection onto **P** and **Q**.

5.3 The Interval Pair

In this section we consider the consistent quantification of intervals by a coordinated set of chains. We will focus on situations where both elements defining the endpoints of the interval both project and dual project onto each of the quantifying chains under consideration. Since coordination is based on the projection of closed intervals situated on one chain to closed intervals on another chain, we begin by examining consistent quantification in that special case.

Figure 8 considers a set of five mutually coordinated chains: **P**, **Q**, **R**, **S**, and **T**, which collectively form a 1+1 dimensional subspace. These are labeled so that the chains **R**, **S**, and **T** are situated between the pair of chains **P** and **Q**, and the chain **T** is situated between the pair of chains **R** and **S**, so that $\mathbf{P}|\mathbf{R}|\mathbf{T}|\mathbf{S}|\mathbf{Q}$. We consider two elements $a, b \in \mathbf{T}$ that form the closed interval $[a, b]_{\mathbf{T}}$. This closed interval both projects and dual projects onto each of the four other quantifying chains in the coordinated set. By mapping the elements a and b onto the other chains, we can obtain a quantification of $[a, b]_{\mathbf{T}}$ based on the following ***4-tuples*** of valuations:

$$\begin{array}{l}
(v_{\mathbf{P}}(Pa), v_{\mathbf{P}}(Pb), v_{\mathbf{P}}(\overline{P}a), v_{\mathbf{P}}(\overline{P}b))_{\mathbf{P}} \\
(v_{\mathbf{R}}(Ra), v_{\mathbf{R}}(Rb), v_{\mathbf{R}}(\overline{R}a), v_{\mathbf{R}}(\overline{R}b)))_{\mathbf{R}} \\
(v_{\mathbf{T}}(a), v_{\mathbf{T}}(b), v_{\mathbf{T}}(a), v_{\mathbf{T}}(b))_{\mathbf{T}} \\
(v_{\mathbf{S}}(Sa), v_{\mathbf{S}}(Sb), v_{\mathbf{S}}(\overline{S}a), v_{\mathbf{S}}(\overline{S}b))_{\mathbf{S}} \\
(v_{\mathbf{Q}}(Qa), v_{\mathbf{Q}}(Qb), v_{\mathbf{Q}}(\overline{Q}a), v_{\mathbf{Q}}(\overline{Q}b))_{\mathbf{Q}},
\end{array}$$

where $Ta = \overline{T}a = a$ and $Tb = \overline{T}b = b$. The remainder of this section is focused on examining relationships among these 4-tuples.

Chain projection maps the interval $[a, b]_{\mathbf{T}}$ to two closed intervals on each of the four other chains, such as $[Pa, Pb]_{\mathbf{P}}$ and $[\overline{P}a, \overline{P}b]_{\mathbf{P}}$ along the chain **P**. Since we have shown that closed intervals on chains possess a unique scalar measure called ***length*** given by the usual lattice distance function (13) defined



Figure 8: (a) An illustration of the consistent quantification of a closed interval $[a,b]_{\mathbf{T}}$ on a chain that is the member of a coordinated set of chains. The interval projects to a pair of intervals on each chain of the set. This enables one to quantify the interval with either four values (4-tuple) or a pair of closed interval lengths, called an interval pair. Note that, in this case, the interval pair consists of identical components. As discussed in Section 5.6, this closed interval is characterized as a pure chain-like interval by the other chains in the coordinated set. (b) A generalized interval also projects to a pair of intervals on each chain. However, its relationship to the quantifying chain determines way in which the interval pair is consistently computed (18). Because the components of the interval pair have like sign, this generalized interval is classified as a chain-like interval (see discussion in Section 5.6).

on the chain itself, this enables us to quantify the interval $[a,b]_{\mathbf{T}}$ with respect to the chain $\mathbf{P}$ by the pair of lengths $v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa)$ and $v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)$ corresponding to the two closed intervals $[Pa, Pb]_{\mathbf{P}}$ and $[\overline{P}a, \overline{P}b]_{\mathbf{P}}$ on $\mathbf{P}$ onto which $[a,b]_{\mathbf{T}}$ projects. This the 4-tuple $(v_{\mathbf{P}}(Pa), v_{\mathbf{P}}(Pb), v_{\mathbf{P}}(\overline{P}a), v_{\mathbf{P}}(\overline{P}b))_{\mathbf{P}}$ can be expressed as a pair of scalars $\big(v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa), v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)\big)_{\mathbf{P}}$, which we call the ***interval pair***.

Since the chains $\mathbf{T}$ and $\mathbf{P}$ are coordinated, we have that the closed interval $[a,b]_{\mathbf{T}}$ projects and dual projects to closed intervals of equal length on $\mathbf{P}$, so that we can define $\Delta p = v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa) = v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)$. This results in a quantification with respect to the chain $\mathbf{P}$ consisting of an interval pair with identical components $(\Delta p, \Delta p)_{\mathbf{P}}$, which we refer to as a ***symmetric pair***. Furthermore, the fact that each pair of chains is coordinated, implies that $\Delta p = \Delta r = \Delta t = \Delta s = \Delta q$, so that each of the chains in the coordinated set quantifies the closed interval with identical interval pairs.

The situation is not quite as straightforward for intervals in general. We begin by considering quantification of a generalized interval with endpoints that are situated within the 1+1 dimensional subspace defined by a set of coordinated quantifying chains. Later, Section 5.7, which is focused on orthogonal subspaces, is motivated by inconsistencies in quantification that arise when one or more endpoints of the interval being quantified are situated outside of this 1+1 dimensional subspace. This results in a more general means of consistent quantification that is discussed in Section 7.

Consider the quantification of a generalized interval by a single chain such that they are both situated within the same 1+1 dimensional subspace. There are two possible cases to consider: (Case 1) the endpoints of the interval are both situated on the same side of the quantifying chain, and (Case 2) the endpoints of the interval are situated on opposite sides of the quantifying chain. The specific example of the quantification of a closed interval situated on one of the chains of the coordinated set fully constrains the solution in Case 1, so that the interval is quantified by the chain $\mathbf{P}$ with the pair

$$\big(v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa), v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)\big)_{\mathbf{P}}.$$

We require that quantification in Case 2 be consistent with the quantification obtained in Case 1. Consider the situation illustrated in Figures 8a and 9a and b, where $\mathbf{P}|\mathbf{T}|\mathbf{Q}$ and $\mathbf{P}|a|\mathbf{T}|\mathbf{Q}$ and $\mathbf{P}|\mathbf{T}|b|\mathbf{Q}$ so that the endpoints of the interval $[a, b]$ are situated on the same side of chains $\mathbf{P}$ and $\mathbf{Q}$, but on opposite sides of $\mathbf{T}$, which we denote by $a|\mathbf{T}|b$. Consistency of quantification by chains of the coordinated set requires that quantification by the chain $\mathbf{T}$ be equivalent to the quantification $\big(v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa), v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)\big)_{\mathbf{P}}$ by the chain $\mathbf{P}$. First consider the forward projections onto the chain $\mathbf{P}$. The situation $\mathbf{P}|a|\mathbf{T}$, which corresponds to Case III of (14), implies that

$$P_a = P\overline{T}a.$$

Similarly, the element b satisfies Case II of (14), which results in

$$P_b = PTb.$$

We then have that

$$v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa) = v_{\mathbf{P}}(PTb) - v_{\mathbf{P}}(P\overline{T}a),$$

and by the coordination condition, $v_{\mathbf{P}}(PTb) - v_{\mathbf{P}}(P\overline{T}a) = v_{\mathbf{T}}(Tb) - v_{\mathbf{T}}(\overline{T}a)$ so that

$$v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa) = v_{\mathbf{T}}(Tb) - v_{\mathbf{T}}(\overline{T}a).$$

Applying the same argument to the backward projections, we have that

$$v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a) = v_{\mathbf{T}}(\overline{T}b) - v_{\mathbf{T}}(Ta),$$

so that in Case 2, one can quantify the interval with the pair $\big(v_{\mathbf{T}}(Tb) - v_{\mathbf{T}}(\overline{T}a), v_{\mathbf{T}}(\overline{T}b) - v_{\mathbf{T}}(Ta)\big)$.

The result is any generalized interval $[a, b]$ situated within a 1+1 dimensional subspace defined by a set of coordinated chains can be consistently quantified by any chain $\mathbf{P}$ in the coordinated set with the interval pair defined by

$$[a,b]|_{\mathbf{P}} = \begin{cases} \big(v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(\overline{P}a), v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(Pa)\big)_{\mathbf{P}} & \text{if } a|\mathbf{P}|b \text{ or } b|\mathbf{P}|a \\ \big(v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa), v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)\big)_{\mathbf{P}} & \text{if } [a,b]|\mathbf{P} \text{ or } \mathbf{P}|[a,b] \end{cases} \tag{18}$$

We can define

$$\Delta p = v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa)$$

$$\Delta \overline{p} = v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)$$

so that we can conveniently write the interval pair in the case where the interval is on one side of the quantifying chain as

$$[a,b]|_{\mathbf{P}} = \big(\Delta p, \Delta \overline{p}\big)_{\mathbf{P}}. \tag{19}$$

In the case where the interval is situated between two chains, such as when $\mathbf{P}|a|\mathbf{Q}$ and $\mathbf{P}|b|\mathbf{Q}$, we can use the fact that chains $\mathbf{P}$ and $\mathbf{Q}$ are coordinated to write

$$v_{\mathbf{Q}}(Qb) - v_{\mathbf{Q}}(Qa) = v_{\mathbf{P}}(\overline{P}b) - v_{\mathbf{P}}(\overline{P}a)$$

so that the interval pair is expressed only in terms of forward projections. The interval scalar obtained by quantifying with respect to two chains in this way is denoted $[a,b]|_{\mathbf{PQ}}$ and given by

$$[a,b]|_{\mathbf{PQ}} = \big(v_{\mathbf{P}}(Pb) - v_{\mathbf{P}}(Pa), v_{\mathbf{Q}}(Qb) - v_{\mathbf{Q}}(Qa))\big)_{\mathbf{PQ}} \tag{20}$$

which is more conveniently written

$$[a,b]|_{\mathbf{PQ}} = \big(\Delta p, \Delta q\big)_{\mathbf{PQ}} \tag{21}$$

where $\Delta q = v_{\mathbf{Q}}(Qb) - v_{\mathbf{Q}}(Qa)$. This is useful since the forward projections reflect information that observers represented by chains $\mathbf{P}$ and $\mathbf{Q}$ may be able to access in a practical situation.

5.4 Quantifying Coordinated Chains

In Section 5.1 we demonstrated that chains belonging to a set of coordinated chains can be ordered according to an ordering relation induced by chain projection, and that this gives rise to the concept of a 1+1 dimensional subspace. We can therefore think of an abstract chain of coordinated chains, such as $\mathbf{O} < \mathbf{A} < \mathbf{B} < \mathbf{C} < \mathbf{D}$ illustrated in Figure 10, and consider quantifying the coordinated chains themselves as well as intervals of chains—just as we have previously done for elements along a chain.

An ***interval of coordinated chains*** can be defined as a set of coordinated chains between and including a pair of chains (endpoints) that belong to the coordinated set of chains. For example, given the coordinated set of chains $\mathbf{O} < \mathbf{A} < \mathbf{B} < \mathbf{C} < \mathbf{D}$, the interval of coordinated chains denoted by $[\![\mathbf{A},\mathbf{C}]\!]$ is given by the set of chains $\{\mathbf{A},\mathbf{B},\mathbf{C}\}$. Two intervals of coordinated chains that share common chains can be joined in a way analogous to closed intervals of events along a chain so that

$$[\![\mathbf{A},\mathbf{C}]\!] = [\![\mathbf{A},\mathbf{B}]\!] \cup [\![\mathbf{B},\mathbf{C}]\!] \tag{22}$$



Figure 9: (a) An illustration of the consistent quantification a generalized interval $[a, b]$ by chains in a coordinated set of chains where the components of the interval pair are of opposite sign and opposite in magnitude. With respect to the coordinated set of chains, this interval is classified as a pure antichain-like interval. (b) In general antichain-like intervals are characterized by interval pairs where the components are of opposite sign.

is defined as the set union of the set of chains. Just as we did with elements along a chain and closed intervals, we can assign scalar valuations to chains in the coordinated set, as well as to intervals of coordinated chains. Since the join operation is associative, the function $\oplus$ combining valuations ψ assigned to intervals of coordinated chains is also associative

$$\alpha \oplus (\beta \oplus \gamma) = (\alpha \oplus \beta) \oplus \gamma, \tag{23}$$

where $\alpha = \psi([\![\mathbf{A}, \mathbf{B}]\!])$, $\beta = \psi([\![\mathbf{B}, \mathbf{C}]\!])$, $\gamma = \psi([\![\mathbf{C}, \mathbf{D}]\!])$. As described in Section 4.1, this functional equation for $\oplus$ is called the associativity equation [1, 23], and its general solution is

$$\alpha \oplus \beta = f^{-1}(f(\alpha) + f(\beta)), \tag{24}$$

where f is an arbitrary invertible function [1, 23]. Without loss of generality, we can perform a regraduation to a valuation $D = f \circ \psi$ so that $D([\![\mathbf{A}, \mathbf{B}]\!]) = f(\psi([\![\mathbf{A}, \mathbf{B}]\!]))$ and the valuations assigned to intervals sharing a single common chain are related by

$$D([\![\mathbf{A}, \mathbf{C}]\!]) = D([\![\mathbf{A}, \mathbf{B}]\!] \cup [\![\mathbf{B}, \mathbf{C}]\!]) = D([\![\mathbf{A}, \mathbf{B}]\!]) + D([\![\mathbf{B}, \mathbf{C}]\!]). \tag{25}$$

We refer to the valuation $D([\![\mathbf{A}, \mathbf{B}]\!])$ as the ***distance*** between chains $\mathbf{A}$ and $\mathbf{B}$. Similarly, as described in Section 4.1 for closed intervals of elements, additivity



Figure 10: We can quantify chains in a coordinated set of chains, just as we did events along a chain. We can define an interval of chains as a set of chains between and including the two chains acting as endpoints, and define the distance to be a scalar quantity assigned to the interval. For example $[\![\mathbf{A}, \mathbf{C}]\!] = \{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$. The act of joining intervals of chains is associative , which leads to associativity of the combinations of distances

of D results in

$$D([\![\mathbf{A}, \mathbf{C}]\!]) = X(\mathbf{C}) - X(\mathbf{A}), \tag{26}$$

where $X(\mathbf{A})$ and $X(\mathbf{C})$ are the scalar valuations, referred to as ***position***, assigned to the chains $\mathbf{A}$ and $\mathbf{C}$.

To maintain consistency with the original quantification of the poset, we require that the distance between two chains in a coordinated set of chains must be a function of the valuations of some set of elements projected onto the two chains. Since there can be no preferred elements, we seek a distance assignment that is a function of the interval pair associated with an arbitrary generalized interval defined by a pair of arbitrary elements belonging to each of the two chains, and that all such arbitrary pairs of elements should yield the same distance measure. That is, given a pair of coordinated chains $\mathbf{P}$ and $\mathbf{Q}$ and arbitrary elements on those chains $p \in \mathbf{P}$ and $q \in \mathbf{Q}$ within the coordinated range, we consider the interval $[p, q]$ and require the distance between the chains $D([\![\mathbf{P}, \mathbf{Q}]\!])$ to be a function of $\Delta p = p - Pq$ and $\Delta q = Qp - q$.

Associativity of joining intervals of coordinated chains is related to associativity of the join of projected closed intervals along a coordinated chain, as illustrated in Figure 10. Since this results in additivity of distances as well as lengths of the projected intervals, this implies that the distance $D([\![\mathbf{P}, \mathbf{Q}]\!])$ must be a linear function of the elements of the interval pair $(\Delta p, \Delta q)_{\mathbf{PQ}}$

$$D([\![\mathbf{P}, \mathbf{Q}]\!]) = a\Delta p + b\Delta q \tag{27}$$

for arbitrary $p \in \mathbf{P}$ and $q \in \mathbf{Q}$. Since these elements are arbitrary, we must obtain the same distance using different elements $p' \in \mathbf{P}$ and $q' \in \mathbf{Q}$ so that

$$a\Delta p + b\Delta q = a\Delta p' + b\Delta q' \tag{28}$$

which can be rewritten as

$$a(p - Pq) + b(Qp - q) = a(p' - Pq') + b(Qp' - q') \tag{29}$$

and rearranged to give

$$a(p - p') + b(q' - q) = a(Pq - Pq') + b(Qp' - Qp). \tag{30}$$

Since the chains **P** and **Q** are coordinated, we have that

$$p' - p = Qp' - Qp, \tag{31}$$

and

$$q' - q = Pq' - Pq. \tag{32}$$

Substituting these into the expression above, we have that

$$a(p - p') + b(q' - q) = a(q - q') + b(p' - p). \tag{33}$$

Since $p - p'$ and $q - q'$ are arbitrary, we have that condition that $a = -b$ so that

$$D([\![\mathbf{P}, \mathbf{Q}]\!]) = C(\Delta p - \Delta q), \tag{34}$$

where C is an arbitrary scale. This valuation is guaranteed to give the same distance from chain **P** to chain **Q** given any $p \in \mathbf{P}$ and $q \in \mathbf{Q}$ within the coordinated range.

5.5 The Symmetric-Antisymmetric Decomposition

In the previous section we found that the distance from chain **P** to chain **Q** is an antisymmetric function of the elements of the interval pair $(\Delta p, \Delta q)$ associated with the interval $[p, q]$ where $p \in \mathbf{P}$ and $q \in \mathbf{Q}$ are arbitrary. Let us now consider an arbitrary closed interval $[p, p']$ on **P** and represent the length as a function of the elements of the interval pair found by projecting this closed interval onto both **P** and **Q**. We have that the length $d([p, p']) = p' - p$ is given by

$$d([p, p']) = a\Delta p + b\Delta q, \tag{35}$$

which can be rewritten

$$d([p, p']) = a(p' - p) + b(Qp' - Qp). \tag{36}$$

Since the chains **P** and **Q** are coordinated, we have that

$$p' - p = Qp' - Qp, \tag{37}$$

so that

$$d([p, p']) = a(p' - p) + b(p' - p). \tag{38}$$

Since $d([p, p']) = p' - p$, we have that $a + b = 1$. Choosing a and b so that they have the same magnitude results in $a = b = 1/2$. Thus the length of a closed interval along a chain can be written as a symmetric function of the elements of the associated interval pair;

$$d([p, p']) = \frac{\Delta p + \Delta q}{2}, \tag{39}$$

whereas the distance from one chain to another is an antisymmetric function of the elements of the associated interval pair

$$D([\![\mathbf{P}, \mathbf{Q}]\!]) = \frac{\Delta p - \Delta q}{2}, \tag{40}$$

where we have conveniently adopted the same scale by setting $C = 1/2$ in (34).

These relationships suggest a useful decomposition, which we call the ***symmetric-antisymmetric decomposition***, where an interval pair $(\Delta p, \Delta q)$ is decomposed into the sum of a symmetric pair and an antisymmetric pair

$$(\Delta p, \Delta q)_{\mathbf{PQ}} = \Big(\frac{\Delta p + \Delta q}{2}, \frac{\Delta p + \Delta q}{2}\Big)_{\mathbf{PQ}} + \Big(\frac{\Delta p - \Delta q}{2}, \frac{\Delta q - \Delta p}{2}\Big)_{\mathbf{PQ}}. \tag{41}$$

In this decomposition, the symmetric component represents projected lengths along the two coordinated chains, and the antisymmetric component represents a component of distance between the elements with respect to the subspace induced by the two coordinated chains. We will examine this decomposition in more detail in Section 5.8 when we look at joining generalized intervals.

5.6 Interval Classes

Given a 1+1 dimensional subspace induced by two coordinated chains $\mathbf{P}$ and $\mathbf{Q}$, the interval pair $\big(\Delta p, \Delta q\big)_{\mathbf{PQ}}$ enables us to identify three equivalence classes of generalized intervals $[a, b]$ situated within the 1+1 dimensional subspace induced by $\mathbf{P}$ and $\mathbf{Q}$ based on whether the two components of the pair are of like sign, opposite sign, or whether one of them is zero.

Closed intervals along a chain coordinated with $\mathbf{P}$ and $\mathbf{Q}$ are quantified by a ***symmetric pair***, such as $(\Delta p, \Delta p)$, where the components of the pair are both of like sign and equal magnitude. Since these belong to the first class where the pair components are of like sign, we say that intervals in this equivalence class are ***chain-like*** (see Figures 8 and 11). Note, however, that since these are generalized intervals defined by two events which may be incomparable, ***not all chain-like intervals form chains***. Chain-like intervals quantified by a chain where each component of the interval pair is of equal magnitude (as in the case of a closed interval quantified by the host chain) are referred to as ***pure chain-like*** as in Figure 8a. Though it should be noted that this latter classification is dependent on the relationship between the interval and the quantifying chain as it is not true in general that an interval characterized as pure chain-like by one chain will be characterized as pure chain-like by another chain.

Similarly, generalized intervals which are quantified by interval pairs with components of opposite sign are called ***antichain-like*** (see Figures 9 and 11). All antichain-like intervals are antichains. In the case where the components of the interval pair are equal in magnitude, but opposite in sign, the interval is referred to as ***pure antichain-like*** and the interval pair is said to be an ***antisymmetric pair*** (see Figure 9a). However, just as in the case of a pure chain-like interval, classification as a pure antichain-like interval is as dependent on the interval as it is the quantifying chain.

Generalized intervals that project to the same element on one of the two chains results in either $\Delta p = 0$ or $\Delta q = 0$. In such cases, one element of the generalized interval projects to the other and then to one of the quantifying chains. For this reason, we call such intervals, ***projection-like*** (see Figure 11).



Figure 11: This figure illustrates three classes of relationships between two elements forming a generalized interval and the observer chains. (Top Left) *Antichain-like* intervals have elements that project in opposite order to the two chains resulting in a pair quantification with opposite signs. (Top Center) *Projection-like* intervals are characterized by the fact that the two elements project to the same element on one of the observer chains so that one element of the quantifying pair is zero. (Top Right) *Chain-like* intervals are defined by elements that project to the observer chains in the same order resulting in a quantifying pair with like signs. (Bottom Left) A purely antichain-like interval is characterized by a quantifying pair that is anti-symmetric as in $(\Delta, -\Delta)$. (Bottom Right) A purely chain-like interval is quantified by a symmetric pair as in (Δ, Δ).

For a set of collinear chains, one can show that the elements of the interval pair for any interval cannot change sign when quantified by another chain in the set. For example, assume that there exists an interval $[a, b]$ that projects to the closed interval $[Pa, Pb]$ on the chain $\mathbf{P}$ and the closed interval $[QPb, QPa]$ on the chain $\mathbf{Q}$ where $Pb > Pa$ and $QPa \geq QPb$. Since, the projection of a onto a chain $\mathbf{P}$ is given by the least element of the chain $\mathbf{P}$ that includes a, we have that $QPa \geq QPb > Pb > Pa > a$ so that Qb is the projection of a onto $\mathbf{Q}$. The result is that the closed interval $[Pa, Pb]$ projects either to $[QPa, QPb]$ or to $[QPb, QPb]$, which results in the element of the interval pair having either the same sign or zero. The result is that chain-like intervals can never be antichain-like intervals and vice versa, although either can be observed to be projection-like.

5.7 Orthogonal Subspaces

In the previous sections, we showed how a pair of coordinated chains induces a 1+1 dimensional subspace in the poset. Elements and chains not included in a given subspace may form subspaces of their own. We begin by considering a particular example that motivates the concept of orthogonal subspaces.

Consider a pair of coordinated chains **P** and **Q**, which form a subspace $\overline{\mathbf{PQ}}$, and a second pair of coordinated chains **R** and **S** not in $\overline{\mathbf{PQ}}$, which form a distinct subspace $\overline{\mathbf{SR}}$. Consider $p \in \mathbf{P}$ and $q \in \mathbf{Q}$ such that they form a pure antichain-like generalized interval $[p, q]$ with respect to $\overline{\mathbf{PQ}}$, which when quantified results in

$$\begin{aligned}[p,q]|_{\mathbf{PQ}} &= (\Delta, -\Delta)_{\mathbf{PQ}} \\ &= (v_{\mathbf{P}}(p) - v_{\mathbf{P}}(Pq), v_{\mathbf{Q}}(q) - v_{\mathbf{P}}(Qp))_{\mathbf{PQ}}.\end{aligned}$$

where the notation $[p,q]|_{\mathbf{PQ}}$ indicates that the generalized interval $[p, q]$ is being quantified with respect to the chains **P** and **Q**, which is also indicated by the subscript on the resulting pair.

Now consider that p and q project to **R** and **S** so that $Rp = Rq$ and $Sp = Sq$ resulting in a quantification of $[p, q]$ with respect to $\overline{\mathbf{RS}}$ equal to

$$\begin{aligned}[p,q]|_{\mathbf{RS}} &= (v_R(Rp) - v_R(Rq), v_S(Sp) - v_S(Sq))_{\mathbf{RS}} \\ &= (0,0)_{\mathbf{RS}}.\end{aligned}$$

This is illustrated in Figure 12A-C. Note also that the situation is similar for the antichain-like interval $[\overline{R}p, \overline{S}p]$ in $\overline{\mathbf{SR}}$ in that it is quantified by $(0,0)_{\mathbf{PQ}}$ with respect to $\overline{\mathbf{PQ}}$. In this case we say that the subspaces $\overline{\mathbf{PQ}}$ and $\overline{\mathbf{SR}}$ are ***orthogonal*** to one another.

Figure 12D illustrates the chains shown in 12(C) along with an additional chain **O**, which has been added so that it is collinear with both $\overline{\mathbf{PQ}}$ and $\overline{\mathbf{RS}}$. We call this a ***geometric view*** where each chain is indicated by a dark circle and they are positioned relative to one another based on the betweeness relations $\mathbf{P}|\mathbf{O}|\mathbf{Q}$ and $\mathbf{R}|\mathbf{O}|\mathbf{S}$. The two independent 1+1 dimensional subspaces when combined results in a 2+1 dimensional subspace where the original ordering relation is supplemented with two ordering relations induced by the two independent collinear sets of chains. The geometric view highlights the two induced dimensions while suppressing the original dimension along the chains.

While, Figure 12C illustrates a motivating example, the general situation is more subtle. While an interval quantified by a pure antisymmetric pair with respect to the chains **P** and **Q** is quantified by $(0, 0)$ with respect to the chains **R** and **S**, that very same interval will not be quantified by $(0, 0)$ with respect to the pair of chains **R** and **O** or with respect to the pair of chains **O** and **S**. In Section 7, we derive a more advanced method of projection of an interval onto a subspace that yields consistent results when quantified with respect to any pair of chains in that subspace.

5.8 Joining Generalized Intervals

In Section 4.1, we introduced the concept of joining closed intervals along a chain. Here we extend this concept to that of joining generalized intervals. Given two generalized intervals that share a common element, such as $[a, b]$ and



Figure 12: (A) A pure anti-symmetric generalized interval $[p, q]$ in the subspace $\overline{\mathbf{PQ}}$. (B) Projection of $[p, q]$ onto $\mathbf{RS}$ is quantified by the pair $(0, 0)_{RS}$. (C) An illustration of the relationship between the two subspaces where pure antichain-like intervals in $\overline{\mathbf{RS}}$ cannot be quantified by $\overline{\mathbf{PQ}}$ vice versa. (D) Illustrates the chains shown in (C) in a ***geometric view*** where each chain is indicated by a dark circle and they are positioned relative to one another based on the betweeness relations. The chain $\mathbf{O}$ has been added so that it is collinear with both $\overline{\mathbf{PQ}}$ and $\overline{\mathbf{RS}}$ and situated between the two chains in each pair.

$[b, c]$, we can define a map $\uplus$ that takes these two intervals to a third unique interval given by

$$[a, c] = [a, b] \uplus [b, c]. \tag{42}$$

In the case where each of these intervals is situated between a pair of coordinated chains $\mathbf{P}$ and $\mathbf{Q}$ in the $\overline{\mathbf{PQ}}$ subspace, the interval $[a, c]$ is quantified by the pair $(p_c - p_a, q_c - q_a)_{\mathbf{PQ}}$, the interval $[a, b]$ by the pair $(p_b - p_a, q_b - q_a)_{\mathbf{PQ}}$, and the interval $[b, c]$ by the pair $(p_c - p_b, q_c - q_b)_{\mathbf{PQ}}$ so that the resulting interval pairs sum in accordance with the $\uplus$ operation

$$(p_c - p_a, q_c - q_a)_{\mathbf{PQ}} = (p_b - p_a, q_b - q_a)_{\mathbf{PQ}} + (p_c - p_b, q_c - q_b)_{\mathbf{PQ}}. \tag{43}$$

This enables us to decompose intervals by introducing artificial events defined by their projection onto a pair of coordinated chains $\mathbf{P}$ and $\mathbf{Q}$. Given two events a and b situated between two coordinated chains $\mathbf{P}$ and $\mathbf{Q}$, we can quantify the interval with the pair $(p_b - p_a, q_b - q_a)$. Consider a decomposition

$$[a, b] = [a, 0] \uplus [0, b] \tag{44}$$

where the pair $(p_0 - p_a, q_0 - q_a)$ quantifying $[a, 0]$ is defined to be antisymmetric so that the two components of the pair are of equal magnitude but opposite sign, $p_0 - p_a = q_a - q_0$, and the pair $(p_b - p_0, q_b - q_0)$ quantifying $[0, b]$ is defined to be symmetric so that the two components of the pair are of equal sign and magnitude $p_b - p_0 = q_b - q_0$. These conditions are satisfied by an artificial event 0 defined by the projections

$$\begin{aligned} p_0 &= \frac{1}{2}(p_a + p_b + q_a - q_b) && (45) \\ q_0 &= \frac{1}{2}(p_a - p_b + q_a + q_b). && (46) \end{aligned}$$

The result is that any interval situated in the subspace defined by two coordinated chains can be expressed in terms of the join of a pure chain-like interval quantified by a symmetric pair and a pure antichain-like interval quantified by an antisymmetric pair in accordance with the symmetric-antisymmetric decomposition (illustrated in Figure 13)

$$(\Delta p, \Delta q)_{\mathbf{PQ}} = \Big(\frac{\Delta p + \Delta q}{2}, \frac{\Delta p + \Delta q}{2}\Big)_{\mathbf{PQ}} + \Big(\frac{\Delta p - \Delta q}{2}, \frac{\Delta q - \Delta p}{2}\Big)_{\mathbf{PQ}}. \quad (47)$$

One can also join intervals situated in distinct subspaces, although additivity of interval pairs does not hold. For example, consider chains $\mathbf{P}$, $\mathbf{O}$, and $\mathbf{R}$ in the situation illustrated in Figure 12D and assume that they are pairwise coordinated. Given elements $p \in \mathbf{P}$, $o \in \mathbf{O}$, and $r \in \mathbf{R}$, we can construct intervals where

$$[p, r] = [p, o] \uplus [o, r]. \quad (48)$$

Quantifying $[p, r]$ with respect to the coordinated pair of chains $\mathbf{P}$ and $\mathbf{R}$, we get the interval pair $(p_r - p, r - r_p)_{\mathbf{PR}}$. Quantifying the other two intervals within their respective subspaces, we obtain $(p_o - p, o - o_p)_{\mathbf{PO}}$ and $(o_r - o, r - r_o)_{\mathbf{OR}}$, where

$$(p_r - p, r - r_p)_{\mathbf{PR}} \neq (p_o - p, o - o_p)_{\mathbf{PO}} + (o_r - o, r - r_o)_{\mathbf{OR}}. \quad (49)$$

Instead, we write

$$(p_r - p, r - r_p)_{\mathbf{PR}} \sim (p_o - p, o - o_p)_{\mathbf{PO}} \oplus (o_r - o, r - r_o)_{\mathbf{OR}} \quad (50)$$

where $p = p_p$, $o = o_o$, $r = r_r$, and the operator $\oplus$, which is defined implicitly through the relation in (48), symbolically indicates a decomposition of the interval pair into distinct subspaces.

5.9 Scalar Quantification of Intervals

Next, we explore scalar measures of intervals, which will later form the foundation for a metric. As described above, closed intervals on a chain can be quantified by a scalar length. To extend this concept of length to generalized intervals, we begin by identifying the constraints imposed on our scalar measure by the special case of a closed interval on one chain that projects and dual projects onto two closed intervals on a second chain.

Consider two chains $\mathbf{S}$ and $\mathbf{P}$ where N successive intervals each of length Δs on the chain $\mathbf{S}$ project onto N successive intervals each of length Δp on $\mathbf{P}$ and dual project onto N successive intervals each of length $\Delta \overline{p}$ as illustrated in Figure 14A. Furthermore, we assume that the two chains reside in the same 1+1D subspace so that in the special case where $\Delta p = \Delta \overline{p}$ we have that the



Figure 13: This figure illustrates the symmetric-antisymmetric decomposition in the context of joining generalized intervals. A generalized interval $[a, b]$ is quantified by the pair $(\Delta p, \Delta q) = (p_b - p_a, q_b - q_a)$. This is equivalent to imagining an event 0 defined by the projections (p_0, q_0) onto $\mathbf{P}$ and $\mathbf{Q}$ such that $\Delta t = p_b - p_0 = q_b - q_0$ and $\Delta x = p_a - p_0 = -(q_a - q_0)$. We can define a join operation, $\uplus$, analogous to that in Appendix I, which concatenates two intervals that share a common element by dropping the common element so that $[a, b] = [a, 0] \uplus [0, b]$, where $[a, 0]$ is an antichain-like interval quantified by an antisymmetric pair $(\Delta x, -\Delta x)$ and $[0, b]$ is a chain-like interval quantified by a symmetric pair $(\Delta t, \Delta t)$.

chains are coordinated and that the lengths of the closed intervals are consistent with $\Delta p = \Delta \overline{p} = \Delta s$. We refer to two such chains as being ***linearly-related***.

Given a single closed interval on $\mathbf{S}$ with length Δs, and its quantification by $\mathbf{P}$ with the pair $(\Delta p, \Delta \overline{p})_{\mathbf{P}}$, we aim to identify a function σ that takes the pair $(\Delta p, \Delta \overline{p})_{\mathbf{P}}$ to Δs so that

$$\sigma(\Delta p, \Delta \overline{p}) = \Delta s. \tag{51}$$

The scalar measures assigned to closed intervals on each of the two chains can be rescaled (change of units) and still remain consistent with additivity under the combination operator $\cup$ discussed in Appendix I. Rescaling by a positive real number α results in

$$\begin{aligned} \sigma(\alpha \Delta p, \alpha \Delta \overline{p}) &= \alpha \Delta s, & (52)\\ &= \alpha \sigma(\Delta p, \Delta \overline{p}). & (53) \end{aligned}$$

This functional equation for σ is of the general form known as the ***homogeneity equation***:

$$F(zx, zy) = z^k F(x, y) \tag{54}$$

whose solution can be written as [1]

$$F(x, y) = \begin{cases} (\sqrt{xy})^k \ h(\frac{x}{y}) & \text{if } xy \neq 0 \\ a x^k & \text{if } x \neq 0, y = 0 \\ b y^k & \text{if } y \neq 0, x = 0 \\ c & \text{if } y = 0, x = 0 \end{cases} \tag{55}$$



Figure 14: (A) N successive intervals each of length Δs on the chain $\mathbf{S}$ forward project onto N successive intervals each of length Δp on $\mathbf{P}$ and back project onto N successive intervals each of length $\Delta\overline{p}$. (B) Projections of the chain-like interval $\mathbf{I}$ onto the two linearly-related chains $\mathbf{S}$ and $\mathbf{P}$.

where $h(\frac{x}{y}) = h(\frac{y}{x})$. In our example, the exponent $k = 1$ and there is a symmetry between x and y, so that $a = b$. When $\Delta s = 0$ we have that $\Delta p = \Delta\overline{p} = 0$ so that $\sigma(0,0) = 0$ giving $c = 0$, so that we can write

$$\sigma(x,y) = \begin{cases} \sqrt{xy}\; h(\frac{x}{y}) & \text{if } xy \neq 0 \\ ax & \text{if } x \neq 0, y = 0 \\ ay & \text{if } y \neq 0, x = 0 \\ 0 & \text{if } y = 0, x = 0 \end{cases} \tag{56}$$

By considering the special case where $\mathbf{P}$ and $\mathbf{S}$ are coordinated so that an interval of length Δs on the chain $\mathbf{S}$ is quantified by $\mathbf{P}$ with the pair $(\Delta s, \Delta s)_{\mathbf{P}}$, we find that

$$\sigma(\Delta s, \Delta s) = \Delta s\; h(\frac{\Delta s}{\Delta s}) = \Delta s, \tag{57}$$

which implies that $h(1) = 1$.

Having obtained a general form for the scalar measure, we consider the constraint imposed by a second special case. Figure 14B illustrates a chain-like interval $\mathbf{I}$ that projects onto two linearly-related finite chains $\mathbf{S}$ and $\mathbf{P}$ such that they are all collinear so that $\mathbf{P}|\mathbf{S}|\mathbf{I}$. The closed interval of length Δs on $\mathbf{S}$ is quantified by the pair $(\Delta p, \Delta\overline{p})_{\mathbf{P}}$ on $\mathbf{P}$. If the interval $\mathbf{I}$ is quantified by the pair $(\alpha\Delta s, \beta\Delta s)_{\mathbf{S}}$ with respect to $\mathbf{S}$, then it is quantified by the pair $(\alpha\Delta p, \beta\Delta\overline{p})_{\mathbf{P}}$ with respect to $\mathbf{P}$. Consistency requires that both chains assign the same scalar to the interval so that

$$\sqrt{(\alpha\Delta s)(\beta\Delta s)}\; h\Big(\frac{\alpha\Delta s}{\beta\Delta s}\Big) = \sqrt{(\alpha\Delta p)(\beta\Delta\overline{p})}\; h\Big(\frac{\alpha\Delta p}{\beta\Delta\overline{p}}\Big), \tag{58}$$

which can be simplified to

$$\Delta s\ h\Big(\frac{\alpha}{\beta}\Big) = \sqrt{\Delta p \Delta \overline{p}}\ h\Big(\frac{\alpha \Delta p}{\beta \Delta \overline{p}}\Big). \tag{59}$$

In the special case examined above where $\alpha = \beta$, we have that

$$\Delta s = \sqrt{\Delta p \Delta \overline{p}}\ h\Big(\frac{\Delta p}{\Delta \overline{p}}\Big). \tag{60}$$

Substituting the expression for Δs above into (59), we find after some simplification that

$$h\Big(\frac{\Delta p}{\Delta \overline{p}}\Big) h\Big(\frac{\alpha}{\beta}\Big) = h\Big(\frac{\alpha}{\beta}\frac{\Delta p}{\Delta \overline{p}}\Big)$$

so that, letting $x = \frac{\Delta p}{\Delta \overline{p}}$ and $y = \frac{\alpha}{\beta}$ we have that

$$h(xy) = h(x)h(y). \tag{61}$$

Since $h(1) = 1$, we can write

$$h\Big(\frac{x}{x}\Big) = 1$$

for non-zero x, which can be factored

$$h(x)h\Big(\frac{1}{x}\Big) = 1.$$

The function h is symmetric in the sense that $h(\frac{x}{y}) = h(\frac{y}{x})$, which implies that $h(\frac{1}{x}) = h(x)$ and

$$h(x)h(x) = 1,$$

so that $h(x) = 1$ or $h(x) = -1$. Equation (60) above gives an expression for Δs, which rules out $h(x) = -1$, so that $h(x) = 1$.

The result is that a chain-like interval quantified by the pair $(\Delta p, \Delta \overline{p})$ can be quantified by a unique scalar consistent among linearly-related chains given by

$$\sigma(\Delta p, \Delta \overline{p}) = \sqrt{\Delta p \Delta \overline{p}}. \tag{62}$$

This special case constrains the functional form of this scalar, so that antichain-like intervals must be quantified similarly, but with $\Delta p \Delta \overline{p} \leq 0$, so that $\sqrt{\Delta p \Delta \overline{p}}$ is imaginary. This suggests some kind of orthogonal relationship between pure chain-like intervals and pure antichain-like intervals, which are characterized by symmetric interval pairs and antisymmetric interval pairs, respectively.

To obtain a real-valued scalar quantification, which is some function $g(\Delta p, \Delta \overline{p})$, that applies to both chain-like and antichain-like intervals, we require that it be consistent with the result $\sigma(\Delta p, \Delta \overline{p}) = \sqrt{\Delta p \Delta \overline{p}}$ so that

$$g(\Delta p, \Delta \overline{p}) = F(\sigma(\Delta p, \Delta \overline{p})), \tag{63}$$

where F is an unknown function to be determined. Consider the chain-like interval quantified by the pair $(\Delta t + \Delta x, \Delta t - \Delta x)$ where $\Delta t > \Delta x$. We then have that

$$g(\Delta t + \Delta x, \Delta t - \Delta x) = F(\sqrt{(\Delta t + \Delta x)(\Delta t - \Delta x)}), \tag{64}$$

which can be rewritten without loss of generality as

$$g(\Delta t + \Delta x, \Delta t - \Delta x) = G((\Delta t)^2 - (\Delta x)^2) \tag{65}$$

where G is an unknown function to be determined.

We now rely on the fact that the joining of orthogonal intervals is associative so that

$$((\Delta x, -\Delta x) \oplus (\Delta y, -\Delta y)) \oplus (\Delta z, -\Delta z) \\ = (\Delta x, -\Delta x) \oplus ((\Delta y, -\Delta y) \oplus (\Delta z, -\Delta z)) \tag{66}$$

where we have dropped the reference to the quantifying chains in each frame. The corresponding scalar quantification is also associative

$$(g(\Delta x, -\Delta x) \hat{\oplus}\, g(\Delta y, -\Delta y)) \hat{\oplus}\, g(\Delta z, -\Delta z) \\ = g(\Delta x, -\Delta x) \hat{\oplus} (g(\Delta y, -\Delta y) \hat{\oplus}\, g(\Delta z, -\Delta z)). \tag{67}$$

where $\hat{\oplus}$ is a function to be determined that combines the scalars of orthogonal intervals. By defining $u = g(\Delta x, -\Delta x)$, $v = g(\Delta y, -\Delta y)$, and $w = g(\Delta z, -\Delta z)$ we can rewrite the equation above as

$$(u \hat{\oplus} v) \hat{\oplus} w = u \hat{\oplus} (v \hat{\oplus} w). \tag{68}$$

This functional equation for $\hat{\oplus}$ is the *associativity equation*, which is the very same functional equation (6) discussed in Appendix 1 within the context of joining closed intervals along a chain. The general solution (7) is given by [1]

$$u \hat{\oplus} v = f(f^{-1}(u) + f^{-1}(v)) \tag{69}$$

where f is an arbitrary invertible function. This indicates that there exists a convenient representation where the scalar measures of orthogonal intervals are additive.

Pure chain-like intervals also enjoy associativity when joined with pure antichain-like intervals so the above theorem applies to them as well. This implies that there exists an additive scalar given by $f^{-1} \circ g$ such that

$$f^{-1}(g(\Delta t + \Delta x, \Delta t - \Delta x)) = f^{-1}(g(\Delta t, \Delta t)) + f^{-1}(g(\Delta x, -\Delta x)) \tag{70}$$

and by applying (65) we have

$$f^{-1}(G((\Delta t)^2 - (\Delta x)^2)) = f^{-1}(g(\Delta t, \Delta t)) + f^{-1}(g(\Delta x, -\Delta x)). \tag{71}$$

Since an interval quantified by the symmetric pair $(\Delta t, \Delta t)$ is chain-like, we have that

$$g(\Delta t, \Delta t) = G((\Delta t)^2), \tag{72}$$

which allows us to obtain an expression for $f^{-1}(g(\Delta x, -\Delta x))$

$$f^{-1}(g(\Delta x, -\Delta x)) = f^{-1}(G((\Delta t)^2 - (\Delta x)^2)) - f^{-1}(G((\Delta t)^2)). \tag{73}$$

Since the left-hand side can only depend on x, we have that $f^{-1} \circ G$ must be linear so that

$$f^{-1}(g(\Delta x, -\Delta x)) = -c(\Delta x)^2 \tag{74}$$

and

$$f^{-1}(g(\Delta t, \Delta t)) = c(\Delta t)^2 \tag{75}$$

where c is a constant which amounts to a freedom to select units. For additivity in (70) to hold in general, we find that the constant $a = 0$ in (56).

We can choose units consistent with those employed by the chain so that $c = 1$. The function $f^{-1} \circ g$ represents a regraduated scalar quantification that is additive in the case of joining orthogonal intervals

$$f^{-1}(g(\Delta p, \Delta \overline{p})) = \Delta p \Delta \overline{p}, \tag{76}$$

which we rename with the composite symbol Δs^2

$$\Delta s^2(\Delta p, \Delta \overline{p}) = \Delta p \Delta \overline{p} \tag{77}$$

to indicate its relationship to the scalar length Δs obtained in the case of closed intervals along the quantifying chain. In the case where the interval being quantified is situated between two coordinated chains $\mathbf{P}$ and $\mathbf{Q}$ we have that $\Delta q = \Delta \overline{p}$ and can write

$$\Delta s^2(\Delta p, \Delta q) = \Delta p \Delta q. \tag{78}$$

We call this scalar quantification of an interval, the ***interval scalar***. It is additive when the intervals being joined are orthogonal, and its square root (or square root of its negation) is additive when the intervals being joined have the same ratio of symmetric and antisymmetric components.

Applying these results to the symmetric-antisymmetric decomposition (41)

$$(\Delta p, \Delta q)_{\mathbf{PQ}} = \Big(\frac{\Delta p + \Delta q}{2}, \frac{\Delta p + \Delta q}{2}\Big)_{\mathbf{PQ}} + \Big(\frac{\Delta p - \Delta q}{2}, \frac{\Delta q - \Delta p}{2}\Big)_{\mathbf{PQ}}, \tag{79}$$

we verify that additivity of the interval scalar holds

$$\Delta s^2 = \Delta p \Delta q = \Big(\frac{\Delta p + \Delta q}{2}\Big)^2 - \Big(\frac{\Delta p - \Delta q}{2}\Big)^2. \tag{80}$$

We refer to the resulting quadratic form on the right-hand side of (80) as the ***Minkowski form***, and stress that Δs^2 does not represent some quantity squared, but instead represents the product of two independent quantities Δp and Δq.

5.10 Pair Transformations

In the previous section, we relied on linearly-related chains to constrain the form of the interval scalar. Here we consider how the interval pair obtained by quantifying with respect to one chain transforms when quantified with respect to another linearly-related chain (Figure 15).

A closed interval on $\mathbf{P}$ with length k is quantified with respect to $\mathbf{P}$ by the symmetric pair $(k, k)_{\mathbf{P}}$. This interval projects to a closed interval of length m on $\mathbf{P'}$ and dual projects to a closed interval of length n on $\mathbf{P'}$ resulting in the pair $(m, n)_{\mathbf{P'}}$. We write the pair transformation in terms of the function L that takes one interval pair to another as

$$L_{\mathbf{P} \to \mathbf{P'}}(k, k)_{\mathbf{P}} = (m, n)_{\mathbf{P'}}. \tag{81}$$



Figure 15: This figure illustrates how the pair quantification of an interval **I** obtained using a chain **P** relates to the pair obtained by quantifying with respect to a linearly-related chain **P′**. The pair transformation can be written in terms of the transformation of a closed interval of length k in **P** and its pair quantification by chain **P′**.

The interval **I** projects onto **P** resulting in the pair $(\Delta p, \Delta\overline{p})_{\mathbf{P}}$, and also onto **P′** resulting in the pair $(\Delta p', \Delta\overline{p'})_{\mathbf{P'}}$ so that

$$L_{\mathbf{P}\to\mathbf{P'}}(\Delta p, \Delta\overline{p})_{\mathbf{P}} = (\Delta p', \Delta\overline{p'})_{\mathbf{P'}}. \tag{82}$$

We can write the lengths of the projected intervals in terms of the length k of the closed interval so that $\Delta p = \alpha k$, $\Delta\overline{p} = \beta k$, and $\Delta p' = \alpha m$, $\Delta\overline{p'} = \beta n$. The pair transformation can then be written as

$$L_{\mathbf{P}\to\mathbf{P'}}(\alpha k, \beta k)_{\mathbf{P}} = (\alpha m, \beta n)_{\mathbf{P'}} \tag{83}$$

indicating that the pair transformation is linear in each argument, as expected since the chains are linearly-related. Constant terms in the linear transformation are zero since intervals of zero length always project to intervals of zero length. Writing this linear transformation in terms of functions of m and n, we find that in general

$$L_{\mathbf{P}\to\mathbf{P'}}(x, y)_{\mathbf{P}} = (x\ f(m,n), y\ g(m,n))_{\mathbf{P'}} \tag{84}$$

where the functions f and g are to be determined.

The interval scalar associated with the closed interval allows us to write $k = \sqrt{mn}$ so that (81) becomes

$$L_{\mathbf{P}\to\mathbf{P'}}(k, k)_{\mathbf{P}} = (\sqrt{mn}\ f(m,n), \sqrt{mn}\ g(m,n))_{\mathbf{P'}} = (m,n)_{\mathbf{P'}}. \tag{85}$$

Equating components we find that

$$m = \sqrt{mn}\ f(m,n) \tag{86}$$

and

$$n = \sqrt{mn}\; g(m,n) \tag{87}$$

so that

$$f(m,n) = g^{-1}(m,n) = \sqrt{\frac{m}{n}}. \tag{88}$$

Note that the fact that f and g are inversely related preserves the interval scalar in such situations. In general, the pair transformation from one quantifying chain to another linearly-related chain is given by

$$\begin{aligned} L_{\mathbf{P}\to\mathbf{P}'}(\Delta p, \Delta\overline{p})_{\mathbf{P}} &= (\Delta p', \Delta\overline{p'})_{\mathbf{P}'} \qquad (89)\\ &= \Big(\Delta p\sqrt{\frac{m}{n}}, \Delta\overline{p}\sqrt{\frac{n}{m}}\Big)_{\mathbf{P}'}. \end{aligned}$$

where m and n are determined from (81).

In the case of quantification by coordinated chains $\mathbf{P}$ and $\mathbf{Q}$, we can write the pair $(\Delta p, \Delta\overline{p})_{\mathbf{P}}$ as $(\Delta p, \Delta q)_{\mathbf{PQ}}$ so that

$$\begin{aligned} L_{\mathbf{P}\to\mathbf{P}'}(\Delta p, \Delta q)_{\mathbf{PQ}} &= (\Delta p', \Delta q')_{\mathbf{P}'\mathbf{Q}'} \qquad (90)\\ &= \Big(\Delta p\sqrt{\frac{m}{n}}, \Delta q\sqrt{\frac{n}{m}}\Big)_{\mathbf{P}'\mathbf{Q}'}. \end{aligned}$$

The fundamental nature of the pair of projections is manifest in the simplicity of this transformation. We observe that this is related to the Bondi k-calculus [5], as well as Kauffman's iterant algebra [14] which treats the transformation as a pair-wise multiplication.

6 The Space-Time Picture

In this section, we introduce a change of variables motivated by the identification of three distinct classes of intervals induced by chain projection. We demonstrate that this change of variables results in a metric analogous to the Minkowski metric and reveals that the pair transformation is analogous to a Lorentz transformation in an analogous space-time.

6.1 Space-Time Coordinates

The symmetric-antisymmetric decomposition suggests a convenient change of variables:

$$\Delta t = \frac{\Delta p + \Delta q}{2} \tag{91}$$

$$\Delta x = \frac{\Delta p - \Delta q}{2} \tag{92}$$

where

$$\Delta p = \Delta t + \Delta x \tag{93}$$

$$\Delta q = \Delta t - \Delta x. \tag{94}$$

With this definition, any interval pair $(\Delta p, \Delta q)$ can be written as

$$(\Delta p, \Delta q) = (\Delta t, \Delta t) + (\Delta x, -\Delta x), \tag{95}$$

where we refer to the two pairs on the right as the time and space components, respectively. Similarly, the interval scalar can be written as

$$\Delta p \Delta q = \Delta t^2 - \Delta x^2 \tag{96}$$

which is analogous to the Minkowski metric of flat space-time.

There are a few important observations to make at this point. First, the interval scalar originates from the product of a pair of quantities and is not some fundamental quantity squared, as is suggested by the usual notation Δs^2. Second, while this derivation suggests that the Minkowski metric may always be employed by a chain to quantify intervals, this does not mean that it is the most convenient description. That is, if the time and space components of a sequence of intervals are constant when quantified with respect to one chain, then, in general, it is not true that the time and space components necessarily will be constant when quantified by another chain. That is, one may always use this metric, but lengths and times may vary.

The time and space coordinates can be applied to the pair transformation (90) as well

$$(\Delta p', \Delta q') = \Big(\Delta p \sqrt{\frac{m}{n}}, \Delta q \sqrt{\frac{n}{m}}\Big). \tag{97}$$

Changing variables to coordinates, mixes the pair resulting in a linear transformation

$$\Big(\Delta t' + \Delta x', \Delta t' - \Delta x'\Big) = \Big((\Delta t + \Delta x)\sqrt{\frac{m}{n}}, (\Delta t - \Delta x)\sqrt{\frac{n}{m}}\Big), \tag{98}$$

which can be represented by a matrix multiplication. Solving for $\Delta t'$ and $\Delta x'$, we find that

$$\Delta t' = \frac{\sqrt{\frac{m}{n}} + \sqrt{\frac{n}{m}}}{2}\Delta t + \frac{\sqrt{\frac{m}{n}} - \sqrt{\frac{n}{m}}}{2}\Delta x \tag{99}$$

$$\Delta x' = \frac{\sqrt{\frac{m}{n}} - \sqrt{\frac{n}{m}}}{2}\Delta t + \frac{\sqrt{\frac{m}{n}} + \sqrt{\frac{n}{m}}}{2}\Delta x \tag{100}$$

By defining

$$\beta = \frac{m-n}{m+n}, \tag{101}$$

we obtain a relation analogous to the ***Lorentz transformation*** in coordinate form

$$\Delta t' = \frac{1}{\sqrt{1-\beta^2}}\Delta t + \frac{-\beta}{\sqrt{1-\beta^2}}\Delta x \tag{102}$$

$$\Delta x' = \frac{-\beta}{\sqrt{1-\beta^2}}\Delta t + \frac{1}{\sqrt{1-\beta^2}}\Delta x, \tag{103}$$

which can be further simplified by defining $\gamma = \frac{1}{\sqrt{1-\beta^2}}$ and writing the linear transformation as a matrix multiplication

$$\begin{bmatrix} \Delta t' \\ \Delta x' \end{bmatrix} = \begin{bmatrix} \gamma & -\beta\gamma \\ -\beta\gamma & \gamma \end{bmatrix} \begin{bmatrix} \Delta t \\ \Delta x \end{bmatrix} \tag{104}$$

These results suggest that time and space can be viewed in terms of a uniquely consistent means of quantifying intervals.

6.2 Motion

The quantity

$$\beta = \frac{m-n}{m+n}, \tag{105}$$

introduced in the derivation above is the relevant quantity that relates two linearly-related chains that project to one another in a constant fashion. Its dependence on the antisymmetric component of the pair results in its antisymmetric behavior when the chains are interchanged. That is, β is antisymmetric in the sense that if chain **P** is related to chain **R** by β, then chain **R** relates to chain **P** by $-\beta$. In the special case where $m = n$, we have that $\beta = 0$. This situation represents two coordinated chains, which in the space-time picture is analogous to two observers at rest with respect to one another.

Moreover, this quantity has the extreme values of $\beta = \pm 1$ which correspond to the cases where $m = 0$ and $n = 0$ indicating that all elements of one chain project onto the same element of the other resulting in an interval scalar of zero. This is a degenerate situation in the sense that one chain can quantify the other, but not vice versa. Such intervals are classified as projection-like, which in the space-time picture are analogous to light-like intervals. Since the interval scalar is invariant among linearly-related chains, we have that if β is extremal with respect to one chain, then it must be extremal with respect to all other linearly-related chains. This is analogous to the experimentally observed fact that the speed of light, which is the maximum speed, is invariant for all inertial frames.

When comparing three or more chains that project to one another in a constant fashion, the values of β describing the relationship between pairs of such chains are related by the familiar velocity addition rule, which can be derived from this point as a standard exercise.

It is important to keep in mind that β represents a relationship between chains in a partially-ordered set. There is no motion in a partially-ordered set—only connectivity. This suggests that physical motion can be interpreted in terms of connectivity, or equivalently that motion is a manifestation of interaction.

6.3 Coordinates and the Pythagorean Decomposition

Given the form of the interval scalar, and the fact that it is additive for orthogonal intervals, we find that this leads immediately to the Pythagorean theorem.

Consider the two orthogonal subspaces defined by the coordinated chains $\mathbf{P}|\mathbf{O}|\mathbf{Q}$ and $\mathbf{R}|\mathbf{O}|\mathbf{S}$ as illustrated in Figure 12. Consider also three events $p \in \mathbf{P}$, $o \in \mathbf{O}$ and $r \in \mathbf{R}$ such that they define three pure antisymmetric intervals $A = [p, o]$, $B = [o, r]$, and $C = [p, r]$ so that

$$[p, r] = [p, o] \uplus [o, r]. \tag{106}$$

and

$$(p_r - p, r - r_p)_{\mathbf{PR}} \sim (p_o - p, o - o_p)_{\mathbf{PO}} \oplus (o_r - o, r - r_o)_{\mathbf{OR}} \tag{107}$$

where $\Delta a = p_o - p = -(o - o_p)$, $\Delta b = o_r - o = -(r - r_o)$, and $\Delta c = p_r - p = -(r - r_p)$ so that we can write

$$(\Delta c, -\Delta c)_{\mathbf{PR}} \sim (\Delta a, -\Delta a)_{\mathbf{PO}} \oplus (\Delta b, -\Delta b)_{\mathbf{OR}}. \tag{108}$$

Since the intervals A and B are orthogonal, the interval scalars sum resulting in

$$\Delta a^2 + \Delta b^2 = \Delta c^2, \tag{109}$$

which is the familiar Pythagorean theorem applied to purely space-like intervals.

This enables one to quantify an interval with respect to an extant set of orthogonal subspaces based on projections of that interval onto the chains in those subspaces thus defining a multidimensional coordinate system. This can be made more abstract by defining imaginary coordinate axes based on imagined, but consistent, projections. This is accomplished by introducing an imaginary decomposition chain $\mathbf{O}'$ where the projections of an interval onto this chain are parameterized by a single parameter θ. As an example, consider an interval $[p_a, r_b]$ defined by $p_a \in \mathbf{P}$ and $r_b \in \mathbf{R}$. This interval is quantified by the pair $(p_a - Pr_b, r_b - Rp_a)_{\mathbf{PR}}$, which can be decomposed into the symmetric and antisymmetric pair

$$(p_a - Pr_b, r_b - Rp_a)_{\mathbf{PR}} = (\Delta t, \Delta t) + (\Delta r, -\Delta r) \tag{110}$$

We can then introduce the chain $\mathbf{O}'$ where, introducing the convenient ***dot-star notation*** where $(\Delta t, \cdot) \doteq (\Delta t, \Delta t)$ and $(\Delta r, *) \doteq (\Delta r, -\Delta r)$, we can write

$$(\Delta t, \cdot) + (\Delta r, *) \sim (\Delta t, \cdot) + (\Delta r \;\, f(\theta), *) + (\Delta r \;\, g(\theta), *) \tag{111}$$

with the condition that $f^2(\theta) + g^2(\theta) = 1$ to preserve the interval scalar. There are many such parameterizations one could choose. For the purpose of illustration, we will choose $f(\theta) = \sin\theta$ and $g(\theta) = \cos\theta$ so that

$$(\Delta t, \cdot) + (\Delta r, *) \;\sim\; (\Delta t, \cdot) + (\Delta r \sin\theta, *) + (\Delta r \cos\theta, *), \tag{112}$$

and

$$\Delta t^2 - \Delta r^2 \;=\; \Delta t^2 - \Delta r^2 \sin^2\theta - \Delta r^2 \cos^2\theta. \tag{113}$$

The antisymmetric pair $(\Delta r \sin\theta, *)$ can be further decomposed by introducing an additional decomposition chain along with an accompanying parameterization ϕ, so that

$$(\Delta r \sin\theta, *) \;\sim\; (\Delta r \sin\theta \, f(\phi), *) + (\Delta r \sin\theta \, g(\phi), *), \tag{114}$$

where again $f^2(\phi) + g^2(\phi) = 1$. Choosing sin and cos as before, and rearranging terms, results in

$$\begin{aligned} &(\Delta t, \cdot) + (\Delta r, *) \sim \\ &\quad (\Delta t, \cdot) + (\Delta r \sin\theta \, \cos\phi, *) + (\Delta r \sin\theta \, \sin\phi, *) + (\Delta r \cos\theta, *), \end{aligned} \tag{115}$$

and

$$\Delta t^2 - \Delta r^2 \;=\; \Delta t^2 - \Delta r^2 \sin^2\theta \, \cos^2\phi - \Delta r^2 \sin^2\theta \, \sin^2\phi - \Delta r^2 \cos^2\theta. \tag{116}$$

This is the familiar representation of the Minkowski metric in spherical coordinates (r, θ, ϕ). A straightforward change of variables to $z = r\cos\theta$ and $\rho = r\sin\theta$ results in a representation of the metric in cylindrical coordinates, and a further change to $x = r\sin\theta\,\cos\phi$ and $y = r\sin\theta\,\sin\phi$ results in a representation of the metric in three-dimensional Cartesian coordinates

$$\Delta t^2 - \Delta r^2 \;=\; \Delta t^2 - \Delta x^2 - \Delta y^2 - \Delta z^2. \tag{117}$$



Figure 16: This figure illustrates the subspace projection of an interval onto a pair of coordinated chains. It is a more advanced projection method in the sense that it provides consistent quantification of the interval by any pair of chains in the coordinated set of chains defining the subspace.

7 Subspace Projection

We have seen that intervals can only be consistently quantified via chain projection if they are situated within the subspace defined by the quantifying chains. Here we introduce a more generally useful projection method that results in consistent quantification of intervals by chains in a subspace when using *any pair* of chains in a coordinated set of chains defining a subspace.

Given an element x and a chain $\mathbf{P}$, which is one of several in a coordinated set of chains, one can compute the distance between the element and the chain as

$$d(x,\mathbf{P}) = \frac{(p - Px) - (\overline{P}p - \overline{P}x)}{2}. \tag{118}$$

We then define the projection of the interval $[x,y]$ onto the subspace $\overline{\mathbf{PQ}}$ in terms of the antisymmetric combination of the squares of the four relevant distances

$$(d(y,\mathbf{P})^2 - d(x,\mathbf{P})^2) - (d(y,\mathbf{Q})^2 - d(x,\mathbf{Q})^2) \tag{119}$$

which can be rewritten as

$$(d(y,\mathbf{P})^2 - d(y,\mathbf{Q})^2) - (d(x,\mathbf{P})^2 - d(x,\mathbf{Q})^2). \tag{120}$$

However, this quantity depends on the distance between the chains $\mathbf{P}$ and $\mathbf{Q}$, $d(\mathbf{P},\mathbf{Q})$, defined in (34). Using the Pythagorean theorem (109) one can show that to obtain a consistent quantification of the interval by any pair of coordinated chains within the subspace, we simply normalize $\Pi_{\mathbf{PQ}}[x,y]$ by twice the distance from $\mathbf{P}$ to $\mathbf{Q}$ resulting in

$$\Pi_{\overline{\mathbf{PQ}}}[x,y] = \frac{(d(y,\mathbf{P})^2 - d(y,\mathbf{Q})^2) - (d(x,\mathbf{P})^2 - d(x,\mathbf{Q})^2)}{2d(\mathbf{P},\mathbf{Q})}. \tag{121}$$

By combining the projections of the interval onto two chains in this way, we have developed a more advanced method of projection, which enables one to quantify any generalized interval consistently with respect to a given subspace. This method, which we call ***subspace projection***, is the poset analogue of the ***inner product*** or ***dot product***.

8 Conclusion

We have considered a simple picture of interactions that focuses only on the process of influence, where all potential details of the interactions are stripped away. In this picture, events represent the boundaries of influence with one event representing the act of influencing and the other event representing the corresponding reaction. This results in a partially-ordered set, or a poset, of events.

Consistent apt quantification of ordered structures, such as posets, by assigning n-tuples of numbers (real or natural) to elements or sets of elements, such as intervals, leads to a representation of the ordered structure that can be useful in describing the structure. In cases where ordered sets (or related algebraic structures) possess symmetries, these symmetries will place constraints on any proposed quantification scheme resulting in numeric constraint equations, which can be identified as laws [16]. This has been demonstrated previously in the case of Boolean, and the more general distributive lattices [19] where the sum and product rules of probability theory emerge as constraint equations [23]. Similarly, consistent apt quantification using pairs of numbers applied to the composition of measurement sequences is constrained by the underlying algebraic relations of combining measurements in series and parallel. In this case, the symmetries of the algebraic relations result in constraint equations that have been shown to be equivalent to the sum and product rules of the Feynman path integral formulation of quantum mechanics [12, 11].

However, the fact that posets lack general symmetries means that previously-developed symmetry-based methods of consistent apt quantification cannot be directly applied to posets in general. Here we have shown that the identification of one or more distinguished chains in a poset induces sufficient symmetry to impose useful constraints on quantification and that these constraints in the case of coordinated chains lead directly to the mathematics of special relativity. Specifically, the interval scalar, which represents the unique quantification of intervals (up to a scale) represents the poset analogue of the Minkowski metric. In addition, quantification of an interval with respect to two linearly-related pairs of chains result in two interval pairs that are related by a pair transformation, which is shown to be equivalent to a Lorentz transformation. This enables one to adopt a space-time perspective of the poset, which focuses on the chain-like and antichain-like symmetries induced by the distinguished chain/s at the expense of the simplicity of the mathematics inherent to the poset picture.

Derivation of the Lorentz transformations from causality and fundamental symmetries is not without precedent. Although most past approaches either assume a Minkowski metric, or at the very least, the existence of a space-time manifold endowed with a metric. Zeeman [29] showed that representing causality as a partial ordering on a Minkowski space forces the Lorentz group. Levy-LeBlond [24] produced another derivation that results in both the Lorentz transformation and the Galilean transformation, which does not rely on the Minskowki metric, but rather on the homogeneity and isotropy of space and causality. Kauffman [14] takes this further by considering the principle of relativity as invariance under linear transformations and derives the Lorentz transformation, Galilean transformation, and rotation as special cases. We had been aware of Kauffman's results and the fact that such symmetries result in linear transformations, but after completing our work we were impressed by the similarities between his iterant coordinates and our interval pair, both of which transform in accordance with Bondi's K-calculus [5]. The common element in each of these approaches is the concept of radar time, which manifests itself in our poset approach via chain projection. There has also been a great deal of work to derive space-time geometry (eg. [13, 25]). Of particular note due to its similarity to the present effort is the work by Ehlers, Pirani and Schild [8] who accomplish this by considering properties of light signals along with the assumption of the existence of a space-time manifold. This enables them to extend their results to general relativity by deriving the properties of a curved space-time that is assumed to exist.

After submission of an early version of this paper to the arXiv [22], we were introduced to the work of D'Ariano [6] who showed how the Lorentz transformations can be derived, in principle, from event-counting performed by an observer within a causal network implemented by a quantum computer. This has since been worked out in more detail and most of the quantum mechanical framework has been abstracted away [7]. D'Ariano's approach is similar in spirit to ours in that causality plays a central role, however it relies on a homogeneous causal network, which is unphysical, and focuses on the Lorentz transformation rather than the Minkowski metric.

In contrast to previous approaches, the approach presented here provides additional insights. The antisymmetry of space arises from the antisymmetry of the projections of the antichain-like intervals. As such, antichain-like intervals can be further decomposed into the the join of orthogonal intervals via the Pythagorean theorem; whereas chain-like intervals enjoy no such decomposition. Another way to look at this is to consider that the distinguished chain, which represents a total order, gives rise to one-dimensional time. Sets of multiple coordinated chains induce a different kind of ordering relation, which gives rise to spatial dimensions, which can be multiple in number. Unfortunately, the symmetries introduced here by a distinguished chain are insufficient to constrain the number of spatial dimensions. This can be demonstrated by construction. Consider N finite chains labeled by C_i where $1 \leq i \leq N$. Let each chain C_i have two events $x_i < y_i$. Let $y_i \geq x_j$ for all $1 \leq i \leq N$ and $1 \leq j \leq N$ so that y_i includes the x events on each of the N chains. By symmetry, the distance between each pair of chains in the set of N chains is equal, and thus is only possible in an $(N-1)$ dimensional space. So the present approach admits one-dimensional time and multi-dimensional space. Physics enters by dictating the allowed connectivity of a such a poset, and this is expected to introduce additional constraints that in the future may prove to limit the number of spatial dimensions to three.

Some may note that the current approach fixes the signature of the Minkowski metric to be $(+,-,-,-)$ rather than the $(-,+,+,+)$ more commonly employed in space-time physics. If treated simply as a metric, the signature is arbitrary. However, here the signature arises from the symmetry and antisymmetry of the chain-like and antichain-like components of the interval pair. The fact that these quantities have deeper significance does away with any arbitrariness of the resulting signature. Moreover, the signature derived here agrees with the signature associated with the decomposition of mass into energy and momentum in particle physics. This is significant as it is possible that these quantities may be analogous to rates of interactions in the poset picture.

Another important insight is the fact that the quantity β, which is the poset analogue of speed, is the relevant quantity to describe two linearly-related pairs of chains. In physics, speed is *assumed* to be a relevant quantity. Here it is recovered from the pair transformation and the identification of symmetric pairs with time and antisymmetric pairs with space. It is from this identification of β with speed that we are led to the realization that there exists a finite maximum speed given by $\beta = \pm 1$. This is exemplified by projection-like intervals where the two events defining the interval endpoints project to the same event on a quantifying chain. In addition, the linearity of the pair transformation ensures that the length of the projected interval is always zero so that an interval representing motion at the maximum speed in one frame represents motion at the maximum speed in all linearly-related frames.

More importantly, in the poset picture one views motion in terms of connectivity, which implies that motion is a manifestation of interaction. Particles don't move in this picture. Instead, they transition via discrete jumps with every interaction. This is a bit different from the usual conception of interaction as force. It suggests that the interactions we have quantified, which appear to generate an emergent space-time, do not necessarily cause forces, but at least in some cases enable motion. Due to the intimate connection with space-time, one is led to hypothesize that these interactions may represent a fundamental process from which gravity arises as some sort of side-effect, and as such one might be able to understand the equivalence between inertial and gravitational masses. Such an idea would not be without precedent since it has been suggested that crystal dislocations result in gravity-like behavior in the analogous space-time experienced by an electron moving in graphene [26]. It is possible that the theory presented here may be have something to say about analogous space-time in special materials, such as graphene by considering interactions between the electrons and the crystal lattice as defining a local space-time.

The space-time metric derived here is the Minkowski metric which represents flat space-time. It is important to note that the critical condition in its derivation is the concept of coordinated chains. Additional interactions that act differentially on the coordinated chains will undoubtedly disrupt the coordination condition thus limiting the property of flatness to shorter finite ranges along the chains. We hypothesize that such disruption may be interpreted in terms of a discrete version of curvature.

The discrete nature of interactions in this picture is reminiscent of quantum mechanics (certainly quantum electrodynamics). It would be interesting if the poset picture, in which space-time emerges, could equally support quantum mechanics. If this were the case, it may provide a foundation for quantum gravity. We have demonstrated that a poset model of a free particle that influences its neighbors, but is not influenced by others, reproduces many aspects of Fermion physics [21]. Moreover, from this poset picture, one can derive the Feynman chessboard model of a particle [10] that leads to the Dirac equation in 1+1 dimensions [21]. We are currently continuing these studies to determine if and how space-time physics is related to quantum mechanics in this poset picture.

Galileo wrote, "Measure that which is measurable and make measurable that which is not so." This is the essence behind the concept of quantification. Fundamental symmetries can place strong constraints on a quantification scheme, and in some cases the resulting constraint equations are analogous to what we call laws. In this sense, such laws truly represent an underlying order. By considering a minimalist picture of a physical phenomenon that retains some subset of the fundamental symmetries, one can begin to explore which symmetries may be responsible for a given set of laws by deriving them through apt consistent quantification. As such, the methodology applied here represents a different approach to fundamental, or foundational, physics.

9 Acknowledgements

Kevin Knuth would like to thank Keith Earle, Ariel Caticha, Seth Chaiken, Adom Giffin, Philip Goyal, Jeffrey Jewell, Carlos Rodríguez, Jeff Scargle, John Skilling and Michael Way for many insightful discussions and comments. He would also like to thank Rockne, Ann and Emily Knuth for their support, and Henry Knuth for suggesting that he 'try using a J'. Newshaw Bahreyni would like to thank Shahram Pourmand for his helpful discussions and Mahshid Zahiri, Mohammad Bahreyni and Shima Bahreyni for their continued support. The authors would also like to thank Giacomo Mauro D'Ariano, Alessandro Tosini, Joshua Choinsky, Oleg Lunin, Margaret May, Cristi Stoica and Patrick O'Keefe for valuable comments that have improved the quality of this work.

References

[1] J. Aczél. *Lectures on Functional Equations and Their Applications.* Academic Press, New York, 1966.

[2] G. Birkhoff. *Lattice Theory.* American Mathematical Society, Providence, 1967.

[3] L. Bombelli, J.-H. J.-H. Lee, D. Meyer, and R. Sorkin. Space-time as a causal set. *Phys. Rev. Lett.*, 59:521–524, 1987.

[4] L. Bombelli and D.A. Meyer. The origin of Lorentzian geometry. *Physics Lett. A*, 141:226–228, 1989.

[5] H. Bondi. *Relativity and Common Sense.* Dover, New York, 1980.

[6] G. M. D'Ariano. On the "principle of the quantumness", the quantumness of relativity, and the computational grand-unification. In A. Y. Khrennikov, editor, *Quantum Theory: Reconsideration of Foundations, 5 ed.*, number 1232 in AIP Conference Proceedings, New York, 2010. American Institute of Physics. arXiv:1001.1088 [quant-ph].

[7] G. M. D'Ariano and A. Tosini. Space-time and special relativity from causal networks. 2010. arXiv:1008.4805 [quant-ph].

[8] J. Ehlers, E. Pirani, and A. Schild. The geometry of free fall and light propagation. In L. ORaifeattaigh, editor, *General Relativity / Papers in Honor of J. L. Synge*, pages 63–84. Clarendon Press, Oxford, 1972.

[9] R. P. Feynman. Space-time approach to non-relativistic quantum mechanics. *Rev. Mod. Phys.*, 20(2):367–387, 1948.

[10] R. P. Feynman and A. R. Hibbs. *Quantum Mechanics and Path Integrals.* McGraw-Hill, New York, 1965.

[11] P. Goyal and K. H. Knuth. Quantum theory and probability theory: their relationship and origin in symmetry. *Symmetry*, 3(2):171–206, 2011.

[12] P. Goyal, K. H. Knuth, and J. Skilling. Origin of complex quantum amplitudes and Feynman's rules. *Phys. Rev. A*, 81:022109, 2010. arXiv:0907.0909 [quant-ph].

[13] S. W. Hawking, King A. R., and McCarthy P. J. A new topology for curved space-time which accommodates the causal, differential and causal structures. *J. Math. Phys.*, 17(2):174181, 1976.

[14] L. H. Kauffman. Transformations in special relativity. *Int. J. Theo. Phys.*, 24(3):223–236, 1985.

[15] D. A. Klain and G.-C. Rota. *Introduction to Geometric Probability.* Cambridge Univ. Press, Cambridge, 1997.

[16] K. H. Knuth. Deriving laws from ordering relations. In G. J. Erickson and Y. Zhai, editors, *Bayesian Inference and Maximum Entropy Methods in Science and Engineering, Jackson Hole WY, USA, August 2003*, number 707 in AIP Conference Proceedings, pages 204–235, New York, 2004. American Institute of Physics. arXiv:physics/0403031v1 [physics.data-an].

[17] K. H. Knuth. Lattice duality: The origin of probability and entropy. *Neurocomputing*, 67C:245–274, 2005.

[18] K. H. Knuth. Valuations on lattices and their application to information theory. In *Proceedings of the 2006 IEEE World Congress on Computational Intelligence (IEEE WCCI 2006), Vancouver, BC, Canada, July 2006.*, 2006.

[19] K. H. Knuth. Measuring on lattices. In P. Goggans and C.-Y. Chan, editors, *Bayesian Inference and Maximum Entropy Methods in Science and Engineering, Oxford, MS, USA, 2009*, AIP Conference Proceedings 1193, pages 132–144, New York, 2009. American Institute of Physics. arXiv:0909.3684v1 [math.GM].

[20] K. H. Knuth. Information physics: The new frontier. In A. Mohammad-Djafari P. Bessiere, J.-F. Bercher, editor, *Bayesian Inference and Maximum Entropy Methods in Science and Engineering, Chamonix, France, 2010*, AIP Conference Proceedings 1305, pages 3–19, New York, 2010. American Institute of Physics. arXiv:1009.5161v1 [math-ph].

[21] K. H. Knuth. Inferences about interactions: Fermions and the Dirac equation. In R. Fischer U. von Toussaint, editor, *Bayesian Inference and Maximum Entropy Methods in Science and Engineering, Garching, Germany, 2012*, AIP Conference Proceedings, New York, 2012. American Institute of Physics.

[22] K. H. Knuth and N. Bahreyni. A derivation of special relativity from causal sets., 2010.

[23] K. H. Knuth and J. Skilling. Foundations of inference. *Axioms*, 1(1):38–73, 2012.

[24] J.-M. Levy-LeBlond. One more derivation of the Lorentz transformation. *Am. J. Phy.*, 44(3):271–277, 1976.

[25] D. B. Malament. The class of continuous timelike curves determines the topology of spacetime. *J. Math. Phys.*, 18:13991404, 1977.

[26] A. Mesaros, D. Sadri, and J. Zaanen. Parallel transport of electrons in graphene parallels gravity. *Phys. Rev. B*, 82:073405, Aug. 2010.

[27] R. D. Sorkin. Causal sets: Discrete gravity. In Gomberoff A. and D. Marolf, editors, *Lectures on Quantum Gravity*, pages 305–327. Springer US, 2005.

[28] R. D. Sorkin. Geometry from order: causal sets. In *Einstein Online Vol. 2*, page 1007. 2006.

[29] E. C. Zeeman. Causality implies the Lorentz group. *J. Math. Phys.*, 5(4):490493, 1964.